

Strasbourg, 6 December 2017
[Inf08e_2017.docx]

T-PVS/Inf (2017) 8

CONVENTION ON THE CONSERVATION OF EUROPEAN WILDLIFE
AND NATURAL HABITATS

Standing Committee

37th meeting
Strasbourg, 5-8 December 2017

CODE OF CONDUCT FOR INVASIVE ALIEN TREES

*Document prepared by
Mr Giuseppe Brundu & Mr David M. Richardson
(Department of Agriculture, University of Sassari, Italy - Centre for Invasion Biology, Department of
Botany & Zoology, Stellenbosch University, South Africa)
on behalf of the Bern Convention*

CODE OF CONDUCT FOR INVASIVE ALIEN TREES

by

Giuseppe Brundu & David M. Richardson

(Department of Agriculture, University of Sassari, Italy - Centre for Invasion Biology,

Department of Botany & Zoology, Stellenbosch University, South Africa)

Summary

1. Rationale and aims of the Code of Conduct	4
2. Code of Conduct	5
2.1 Focus of the Code of Conduct: Invasive Alien Trees	5
2.2 Audience and aims of the Code of Conduct.....	6
2.3 A voluntary tool	6
2.4 Implementing, monitoring and evaluating the Code of Conduct	6
3. The principles of the Code of Conduct	6
3.1 Awareness.....	6
3.1.1 Be aware of regulations concerning invasive alien trees	6
3.1.2 Develop systems for information sharing and training programmes	9
3.2 Prevention & Containment	9
3.2.1 Promote – where possible – the use of native trees	9
3.2.2 Adopt good nursery and management practices	10
3.2.3 Adopt good practices for harvesting and transport of timber	11
3.2.4 Adopt good practices for habitat restoration	11
3.3 Early Detection & Rapid Response	12
3.3.1 Promote and implement early detection & rapid response programmes	12
3.3.2 Establish or join a network of sentinel sites	12
3.4 Outreach.....	13
3.4.1 Engage with the public on the risks posed by invasive alien trees, their impacts and on options for management	13
3.5 Forward Planning.....	14
3.5.1 Consider developing research activities on invasive alien trees species and becoming involved in collaborative research projects at national and regional levels.....	14
3.5.2 Take global change trends into consideration	14
4. Background information: a knowledge basis for the Code of Conduct	15
4.1 Benefits arising from planted forests and planted forests of alien trees	15
4.2 Alien tree species in Planted Forests and for other uses: historical and recent pathways of introduction ..	17
4.2.1 General aspects.....	17
4.2.2 Alien conifers	21
4.2.3 Alien eucalypts.....	22

4.2.4 Alien acacias	22
4.2.5 Alien poplars and willows	23
4.2.6 Alien trees and plantations on disturbed land	24
4.2.7 Alien trees in Short-Rotation Forestry, Short-Rotation Coppice.....	24
4.2.8 Alien trees in agroforestry	25
4.2.9 Alien trees in Mediterranean planted forests and sand dune stabilisation	25
4.2.10 Alien trees in arid zones: preventing and combating desertification	26
4.2.11 Genetically improved and genetically modified alien trees.....	27
4.3 Invasive alien trees.....	28
4.3.1 Generalities and key examples	28
4.4 International initiatives and legislation on invasive alien species and invasive alien trees	33
4.4.1 The Convention on Biological Diversity	33
4.4.2 The Council of Europe and the Bern Convention	34
4.4.3 The International Plant Protection Convention	35
4.4.4 The European and Mediterranean Plant Protection Organisation (EPPO)	35
4.5 European initiatives and legislation.....	41
4.5.1 Habitat Directive – Natura 2000.....	41
4.5.2 The Plant Health Regime in the European Union	42
4.5.3 The Biodiversity Strategy of the European Union	42
4.5.4 The EU Regulation on invasive alien species	43
4.5.5 Forest policies in the European Union	43
4.5.6 The EU Energy Policy.....	45
5. References	46
6. Annexes.....	71
6.1 Definitions – Glossary	71
6.2 Alien tree species listed in European countries.....	75

1. RATIONALE AND AIMS OF THE CODE OF CONDUCT

This Code of Conduct is addressed to all relevant stakeholders and decision makers in the 47 Member States of the Council of Europe. It is intended to provide guidance to reduce the negative impacts that might originate from an unregulated use and spread of **invasive¹ alien trees**, i.e. those alien tree species whose introduction or spread has been found to threaten or adversely impact upon biodiversity and related ecosystem services.

Alien trees and well-managed planted forests of alien tree species can be useful in providing various forest goods and services and helping to reduce the pressure on natural forests (FAO 2015b) or provide opportunities for adaptation to climate change and global change.

However, a small number of alien trees are invasive or might become invasive – i.e. they spread from planting sites into adjoining areas, and sometimes cause substantial damage upon biodiversity and related ecosystem services. The challenge is to manage alien trees and existing and future planted forests of alien trees to maximize current benefits and opportunities, while minimising risks and negative impacts, without compromising future benefits and land uses.

To this aim, eleven principles are proposed in the present Code of Conduct:

- Be aware of regulations concerning invasive alien trees;
- Be aware of which alien tree species are invasive or that have a high risk of becoming invasive, and of the invasion debt;
- Develop systems for information sharing and training programmes;
- Promote – where possible – the use of native or non invasive alien trees
- Adopt good nursery and management practices;
- Adopt good practices for habitat restoration;
- Promote and implement early detection & rapid response programmes;
- Establish or join a network of sentinel sites;
- Engage with the public on the risks posed by invasive alien trees, their impacts and on options for management;
- Consider developing research activities on invasive alien trees species and becoming involved in collaborative research projects at national and regional levels;
- Take global change trends into consideration as changing climatic conditions may lead to changes in the species' composition of forests.

¹ Cf. the following section 2.1 for the definition of “invasive alien tree used in the present Code.

2. CODE OF CONDUCT

2.1 Focus of the Code of Conduct: Invasive Alien Trees

In accordance with the CBD² definition, the term **alien**³ **tree** has exclusively a biogeographical meaning, i.e. it refers to a tree species, subspecies or lower taxon, introduced outside its natural past or present distribution and it includes any part, seeds, or propagules of such species that might survive and subsequently reproduce. As such, the term alien tree does not include any negative evaluation of the tree species.

Alien trees can be present outside their native range in confined environments (e.g. Botanic Gardens and Arboreta, plantations) or can be established in the environment. In this Code, the term **naturalised**⁴ **alien tree** species will be used to indicate those alien tree species that have self-sustaining populations, undergo natural dispersal and become incorporated within the resident flora in the environment.

In accordance with the CBD definition, and for the purposes of this Code, an **invasive alien tree** is herewith defined as an alien tree species whose introduction and/or spread threaten or adversely impact upon biodiversity and related ecosystem services.

Not all alien trees will become naturalised. Similarly, **invasive alien trees** are a subset of the naturalised alien trees, as many naturalised alien trees do not go on to become invasive, i.e. they do not threaten biological diversity and do not cause economic or environmental harm.

Therefore, the terms *alien tree*, *naturalised alien tree* and *invasive alien tree* are used throughout this Code of Conduct, with the different meaning as above explained, in accordance with the CBD definitions (COP 6 Decision VI/23), the Recommendations of the Standing Committee (Convention on the Conservation of European Wildlife and Natural Habitats) and the Regulation (EU) No. 1143/2014. The terms *planted forest* is used in accordance with the FAO definitions (FAO 2012, 2015a, 2015b).

² (Decision V/8 of the Conference of the Parties to the Convention on Biological Diversity).

³ In the context of the present Code of Conduct the terms alien, non-native, exotic and introduced tree are considered as equivalent. In accordance with the CBD definition, the term alien tree has exclusively a biogeographical meaning, i.e. it refers to a species, subspecies or lower taxon, introduced outside its natural past or present distribution; it includes any part, seeds, or propagules of such species that might survive and subsequently reproduce. As such, the term alien tree does not include any negative evaluation of the tree species. Only a small percentage of all the alien tree species are, or may become after some time, invasive alien tree species (COP 6 Decision VI/23 “Alien species that threaten ecosystems, habitats or species”). Importantly, an alien tree species is “introduced outside its natural past or present distribution” deliberately or accidentally by man. The definition takes into consideration the Recommendation No. 142 (2009) of the Standing Committee (Convention on the Conservation of European Wildlife and Natural Habitats), adopted on 26 November 2009, interpreting the CBD definition of invasive alien species to take into account climate change, “recommends Contracting Parties to the Convention and invites Observer States to: 1. interpret the term “alien species” for the purpose of the implementation of the European Strategy on Invasive Alien Species as **not including native species naturally extending their range in response to climate change**” (Cf. Section 4.6.2 in this Code). As a result, also past mass migratory events in forest tree populations, postglacial recolonization routes and similar events are not considered herewith in the definition of alien tree species. We focus on alien trees deliberately or accidentally introduced by man outside its natural past or present distribution, where “past” refers to the definition of “neophytes” (i.e. introduced after the 1,500) as used in the CBD context and defined by Pyšek et al. (2004). In addition, with specific concern to this Code, an alien tree species is alien to all the 47 Member States of the Council of Europe. According to this definition the term ‘alien tree species’ does not include foreign provenances of tree species that are native in at least one of the 47 Member States of the Council of Europe.

⁴ According to Regulation EU no. 1143/2014 (Art. 3), an invasive alien species is in the naturalisation stage when its population is self-sustaining.

2.2 Audience and aims of the Code of Conduct

This Code of Conduct is addressed to all relevant stakeholders and decision makers in the 47 Member States of the Council of Europe. It aims to enlist the co-operation of the Forest sector (trade and industry, national forest Authorities, certification bodies and environmental organizations) and associated professionals in preventing, reducing and controlling possible risks and negative impacts. In particular, it is intended to reduce the risk of the introduction of new invasive alien tree species and the negative impacts that might originate from the unregulated use of invasive alien trees. Containment of invasive alien trees to areas set aside for their cultivation or use must become an integral part of management.

It complements the Code of Conduct on Horticulture and Invasive Alien Plants published by the Council of Europe (Heywood & Brunel 2009, 2011) aimed at the horticultural industry and trade and the European Code of Conduct for Botanic Gardens on Invasive Alien Species (Heywood & Sharrock 2013).

2.3 A voluntary tool

This Code of Conduct is voluntary.

This Code does not replace any statutory requirements under international or national legislation but should be seen as complementary to them. Although voluntary, it is important that as many stakeholders as possible should adopt the good practices outlined in this Code so as to reduce the likelihood of compulsory legislation having to be introduced should self-regulation fail. Private forest enterprises, local authorities and public forest managers may wish to publicize their adherence to the Code through adopting a symbol or logo indicating this.

2.4 Implementing, monitoring and evaluating the Code of Conduct

To be fully effective and to increase the likelihood of a long-term behaviour change, a voluntary Code should be widely disseminated and translated into national languages. This clearly stresses the importance of information campaigns aimed at preventing lack of knowledge, possibly coordinated by the key stakeholder's associations and with the support of the national authorities. A straightforward example is provided for by the implementation of the Code of conduct on invasive alien plants in Belgium during the AlterIAS LIFE+ project (Halford et al. 2014). National authorities should acknowledge that the issue of invasive alien trees is a major threat for species, habitats and ecosystems, and undertake measures to ensure that all the available legislation established to prevent introductions of invasive species from Forestry is fully understood, and effectively transposed, implemented and enforced.

National authorities should develop strategies and protocols for dealing objectively with conflicts of interest between those who benefit from the introduction, dissemination and cultivation of invasive alien trees, and those who perceive, and are affected by, negative impacts of these invasive alien trees.

3. THE PRINCIPLES OF THE CODE OF CONDUCT

The eleven principles of this Code of Conduct are clustered in five groups: (1) Awareness; (2) Prevention & Containment; (3) Early Detection & Rapid Response; (4) Outreach and (5) Forward Planning.

3.1 Awareness

3.1.1 *Be aware of regulations concerning invasive alien trees*

Those engaged in the introduction of new tree species and in the planted forest sector need to be aware of their obligations under regulations and legislation. The main obligations under existing laws and treaties are detailed in the Section 4 of this Code.

The Regulation (EU) No. 1143/2014, the Plant Health Directive 2000/29/EC, the Wildlife Trade Regulations (338/97/EC and 1808/2001/EC) and the Habitats Directive (92/43/EEC) only apply to the member countries of the European Union. Many other international conventions addressing issues of

invasive alien species have been ratified by European and Mediterranean Countries (Shine 2007). These recommendations may be implemented in the European Union or in national legislation (of countries that ratified these treaties) and lead to the regulation on harmful organisms, inspections, phytosanitary measures, possession, trade and release in the wild of invasive alien plants and quarantine pests. These regulations have an impact on the everyday work in the planted forest sector and in the use of invasive alien trees.

At the national (or subnational) level, some countries have legislation and/or regulations aimed at preventing possession, transport, trade or release in the wild of specific invasive alien trees⁵.

3.1.2 *Be aware of which alien tree species are invasive or that have a high risk of becoming invasive, and of the invasion debt*

Increasing awareness of problems associated with invasive alien trees means that information on invasive alien tree species and ways of dealing with them is becoming more easily accessible (e.g., European Union dedicated WEB site⁶, EPPO⁷, EASIN⁸). There is strong evidence that invasive alien tree species may replicate invasive behaviour in environmentally-similar conditions in different parts of the world.

Invasive alien tree species can have negative impacts even when they are not fully established or widespread (i.e., before they become naturalised sensu FAO FRA 2015)⁹.

The fact that some invasive alien forestry trees have not yet spread from given planted forests should not be taken as evidence that spread and negative impacts will not occur in the future. Experience with the same invasive alien tree in planted forests in other parts of the world, including areas where the species have long residence times, should be evaluated to assess the extent of “invasion debt” (Richardson et al. 2015)¹⁰.

⁵ For example, in Norway, the 2005 white paper on the Government's environmental policy and the state of the environment in Norway (Report No. 21 - 2004–2005 - to the Storting), the new Forestry Act (Act of 27 May 2005, no. 31, relating to forestry), the Nature Diversity Act (Act of 16 June 2009, no. 100), the Regulation on non-native trees (Regulation of 15 March 2013, no. 284), the national Strategy on Invasive Alien Species (published in May 2007) and the Norwegian Black List (Gederaas et al. 2012), are the main national specific documents referring to non-native trees. The Guidelines on trees, shrubs and plants for planting and landscaping in the Maltese Islands limit the use of alien trees in afforestation projects on agricultural land (MEPA 2002). The Iceland Forest Service has put forth a set of guidelines to afforestation planners: planting of alien trees within natural woodlands is discouraged (Gunnarsson et al. 2005). Planting in treeless land must be carefully assessed considering the phenomenal and unique importance of the Icelandic breeding waterfowl populations which are at risk from the forestry (*Cf.* Bern Convention, Recommendation No. 96 (2002) on conservation of natural habitats and wildlife, especially birds, in afforestation of lowland in Iceland (<https://wcd.coe.int/>). The Swedish Forestry Act placed restrictions on the planting programme of *P. contorta* in 1987, 1989 and 1991 due to extensive infection by *Gremmeniella abietina* in high elevation areas in northern Sweden after periods of extreme weather conditions from 1984 to 1987 (Karlman 2001). Many countries have national or sub-national “black lists” or other types of lists, identifying those alien tree species whose introduction is prohibited or discouraged due to their potential adverse effects on the environment or human, animal or plant health. An alternative approach used in other countries relies on the “white list” (or red, green and amber lists, see Perrings et al. 2005; Simberloff 2006) of low invasion risk alien species, including trees.

⁶ http://ec.europa.eu/environment/nature/invasivealien/list/index_en.htm

⁷ https://www.eppo.int/INVASIVE_PLANTS/ias_plants.htm

⁸ <https://easin.jrc.ec.europa.eu/>

⁹ Some tree species can have negative impacts for example, allergic pollen can affect human health, they can act as vectors of new pests or pathogens for other forest tree species (e.g., Engelmark et al. 2001), they can modify ground vegetation, soil properties and soil fauna (Finch & Szumelda 2007), water balance, fire resilience at the stand level, .

¹⁰ According to Richardson et al. (2015) invasion debt is composed by four main components: (1) the number of species not yet introduced but likely to be introduced in the future given current levels of introduction/propagule pressure; (2) the establishment of introduced species; (3) the potential increase in area invaded by established species

Invasive alien tree species included in “black-lists” should not be released in the environment or used for new planted forests in the countries or regions where they are listed. Additionally, in a huge country the translocation of a native tree species from one part to another has to be carefully assessed to take into account both opportunity and risks. For this reason, for Russia, Notov et al. (2011) propose the adoption of three-level system of sub-national lists called “black books”.

Nevertheless, lists offer a positive approach for both companies and government agencies and could be used to fast-track approval of or reduce liability for forest owners when using low-risk non-native trees for plantations.

For each new alien tree species or provenance¹¹ introduced which has not already been evaluated, those¹² introducing the new alien tree species or planning new planted forests with the new alien tree species should run a risk assessment or risk analysis to take into account both opportunity and risks. According to Křivánek & Pyšek (2006)¹³ two main groups of risk assessment models can be considered, based on the methods used and the phase of the invasion process they target: (1) pre-introduction models predicting the potential behaviour of an alien tree species prior to its introduction; (2) post-introduction models predicting the future behaviour of an alien tree species that have already become naturalized or invasive in the new area. There are over 100 risk assessment and risk analysis schemes for plant species (Leung et al. 2012; Křivánek & Pyšek (2006), with some decision schemes developed specifically for trees or woody plants (Reichard and Hamilton 1997; Pheloung et al. 1999; Haysom & Murphy 2003; Widrlechner et al. 2004; Kumschick & Richardson 2013; Wilson et al. 2014). In 2014 only a few risk assessment methods were in line with the requirements of the Regulation (EU) No. 1143/2014 (Roy et al.

(including invasive species); (4) and the potential increase in impacts. These Authors suggest that invasion debt is a valuable metric for reporting on the threats attributable to biological invasions, that invasion debt must be factored into strategic plans for managing global change, and, as with other studies, they highlight the value of proactive management. However, given the uncertainty associated with biological invasions, further work is required to quantify the different components of invasion debt.

¹¹ “Provenance” in forestry science refers to the particular place where trees are growing or the place of origin of seeds or trees. For example, Norway spruce from different European countries, especially from Germany and Austria, has been used in afforestation in Norway for several decades. Such foreign provenances may differ in adapted ecological traits, such as phenology, frost hardiness, production and spread of seeds, resulting in different growth- and spread potential of the provenances (Aarrestad et al. 2014). In fact, most plant species exhibit spatial structuring of genetic variation throughout their range (Hamrick 1990). While translocation of individual plant species is sometimes proposed as a strategy to increase genetic variation within populations, individuals transferred across different environments may be poorly adapted to the new conditions. Furthermore, there is a risk of outbreeding depression or genetic swamping when divergent populations interbreed. These consequences of seed transfer must therefore be weighed against the potential benefits of increased genetic variation within founding populations (O’Brien et al. 2007; Aitken & Bemmels 2016). Cf. also the Council Directive 1999/105/EC of 22 December 1999 on the marketing of forest reproductive material (Article 2 (g): “for a species or sub-species, the region of provenance is the area or group of areas subject to sufficiently uniform ecological conditions in which stands or seed sources showing similar phenotypic or genetic characters are found, taking into account altitudinal boundaries where appropriate.”).

¹² Various authors suggest that importers, developers and growers who are responsible for introducing potentially invasive alien species should be responsible for damages to the environment (i.e., ‘polluter pays’ principle), rather than allowing that burden to be borne by tax payers or neighbouring private landowners who are affected (Richardson 1998a; Buddenhagen et al. 2009; Chimera et al. 2010; Witt 2010; McCormick & Howard 2013; Lorentz & Minogue 2015).

¹³ To assess the validity of previously developed risk assessment schemes in the conditions of Central Europe, Křivánek & Pyšek (2006) tested the (1) Australian weed risk assessment scheme (A-WRA; Pheloung et al. 1999); (2) the A-WRA with additional analysis by Daehler et al. (2004); and (3) the decision tree scheme of Reichard & Hamilton (1997) developed in North America, on a data set of 180 alien woody species commonly planted in the Czech Republic. The study revealed that the A-WRA model, especially with additional analysis, appears to be a promising template for building a widely applicable system for screening out invasive plant introductions in the central European region. Gordon et al. (2011, 2012) used the A-WRA to evaluate 38 commercially important *Eucalyptus* species.

2014)¹⁴, however a lot of progress has been made meanwhile and risk assessment protocols have made efforts to comply with the regulation, see also an overview of risk assessments published by Roy et al .2017.

3.1.3 *Develop systems for information sharing and training programmes*

The efficacy of any strategy to address invasive alien trees, including the capacity to produce reliable risk assessment reports (see principle 3.1.2), depends on the available information, and on the sharing of data, knowledge and experience¹⁵. Information sharing systems would greatly improve the ability of authorities to prevent the introduction and spread of invasive alien tree species (Katsanevakis et al. 2014). Also, invasive alien tree species management requires specialist knowledge and skills which can only be developed over time. The capacity and awareness of land owners, forestry officials and other stakeholders are crucial for the effective implementation of the principles of this Code.

3.2 Prevention & Containment

Actions aiming at preventing the potential risk posed by invasive alien trees might often be very useful also to contrast or limit the spread of other alien species in general. It is necessary to take actions aiming to prevent potential risks posed by invasive alien trees; below are some of the key approaches to the matter.

3.2.1 *Promote – where possible – the use of native trees*

The use of native species or non-invasive alien or less-invasive¹⁶ alien tree species as alternatives for highly invasive alien tree species should be always considered¹⁷, wherever possible, as should the precise provenance¹⁸ of seeds and germplasm¹⁹.

¹⁴ The article 5 of the Regulation (EU) No. 1143/2014 list the criteria required to perform a risk assessment for the purposes of article 4, i.e. for the procedure to follow to identify the invasive alien species of Union Concern. Other risk assessment schemes, with additional or different criteria, shall be carried out in relation to the current and potential range of the invasive alien species to be assessed, for other specific purposes.

¹⁵ See also COP 6 Decision VI/23 “Alien species that threaten ecosystems, habitats or species”, Guiding principle 8: Exchange of information.

¹⁶ In France and Belgium (Biogeco, INRA, Univ. Bordeaux, Gembloux Agro-Bio Tech, Université de Liège, Gembloux and BFP, INRA, Univ. Bordeaux, Villenave d'Ornon) a multidisciplinary study on *Robinia pseudoacacia* genetic and phenotypic diversity is evaluating the difference in behaviour of existing populations of this species (http://www.neobiota2016.org/wp/wp-content/uploads/neobiota_2016_book-of-abstracts_web.pdf).

¹⁷ FAO Principle 9 “Conservation of biological diversity” states that “...FAO encourages the establishment of planted forests with indigenous species over exotic species, as they produce a wider range of products and benefits, among them a lower environmental risk and an increase in biodiversity. Introduced species should be selected only in relation to specific management objectives, market conditions and ecological site conditions. The decision to plant introduced species should carefully evaluate the risk that these species may become invasive and have adverse effects on the local biodiversity...” (FAO 2010c). In addition, FAO Principle 10 “Conservation of biological diversity” states that “... Guidelines include but are not limited to: ... selecting indigenous species for the establishment of planted forests if they are equal to or better than introduced species for the purpose intended...” (FAO 2006b). See also the “Protocol for species introductions” proposed by Haysom and Murphy (2003). Cf. also Richardson (1998); FAO (2010c); Gordon et al. (2012); Lorenz & Minogue (2015); Peltzer et al. (2015).

¹⁸ Cf. Principle 3.2.1 (footnote n. 10).

¹⁹ Cf. the Pan-European guidelines for afforestation and reforestation with a special focus on provisions of the UNFCCC. Adopted by the MCPFE Expert Level Meeting on 12-13 November 2008 and by the PEBLDS Bureau on behalf of the PEBLDS Council on 4 November, 2008. In particular, at Ecological Guidelines, p. 21 “species, provenances, varieties or ecotypes outside their natural range should only be used where their introduction would not endanger important and/or valuable indigenous ecosystems, flora and fauna. Those that are likely to be invasive should be avoided using the CBD Guiding Principles for the Prevention, Introduction, and Mitigation of Impacts of Alien Species that threaten Ecosystems, Habitats or Species”. Cf. also Aarrestad et al. (2014).

3.2.2 Adopt good nursery and management practices

Best practice methods relating to species and provenances of seed (Karlman 2001), seedling production, weed, pest and disease control should be adopted (FAO 2011). Weeds should be identified, recorded, and eradicated where possible, before planting. The EPPO standard PP 1/141 (3) describes the conduct of trials for the efficacy evaluation of herbicides in tree and shrub nurseries including nurseries within forest stands (EPPO 2009)²⁰.

Nurseries can act as important sources of alien species into planted forest sites. Many forest pests, both insects and pathogens, have also entered new lands via nursery stock²¹.

BOX 3.2.2.1 Sudden oak death (*Phytophthora ramorum*).

Phytophthora ramorum emerged in the US as a forest pathogen causing mortality in oak (*Quercus* spp.) and tanoak (*Notholithocarpus densiflorus*) in California in the mid-1990s, and appeared about the same time in Europe as a nursery pathogen. The pathogen produces spores on a wide variety of foliar hosts, including many popular landscape species. Population genetics studies indicate separate origins for the North American and European populations, and that the North American forest infestation likely originated in nurseries (Ivors et al. 2006; Mascheretti et al. 2008). Although nursery stock has been the major pathway for long-distance spread, the pathogen spreads locally in rain, as well as via surface water runoff from infested nurseries. The pathogen has spread to forests in 14 counties in coastal California and one county in southwest Oregon. In Europe, the pathogen has spread to woodlands in Ireland, the UK, Norway, the Netherlands, and Germany, and has been found in nurseries in sixteen other European countries and Canada (Liebhold et al. 2012). According to Ivors et al. (2006) higher genotypic diversity of *Phytophthora ramorum* in nurseries could be explained by the repeated exchange of pathogen genotypes through the trade of infected plant material, by strong selection pressure selecting new genotypes created through mitotic recombination or mutation, or from both mechanisms. Cultural practices and chemical treatments may be partially responsible for such selection pressure in nurseries. The potential role of plant trade in the creation of an “artificial” panmictic population at the continental level is highlighted by (i) the observation that rare genotypes were found more than once within Europe, particularly in the UK, where the EU4 genotype was found multiple times in different regions, and (ii) the detection of an EU genotype within Oregon and Washington nurseries (Ivors et al. 2006).

Best management practices include criteria such as that biodiversity issues must be considered in planted forest design (CBD COP 11 Decision XI/19, 8 - 19 October 2012 - Hyderabad, India), the design of the planted forest with invasive alien tree shape should to minimise edges at right angles to prevailing winds during seed release season. They should encourage the establishment of representative natural forest within the planted forest and, where possible, restore natural forests on appropriate sites (Secretariat of the Convention on Biological Diversity 2009) and to prevent – whenever possible - plantings of invasive alien trees near “Natura 2000” sites and other protected areas or endangered habitats. Good planted forest practices could also limit the spread of pathogens and pests within planted forests and from infested sites to native species and ecosystems (e.g., Engelmark et al. 2001; FAO 2011).

²⁰ Cf. also, EPPO (2012). EPPO Technical Document No. 1061, EPPO Study on the Risk of Imports of Plants for Planting EPPO Paris. www.eppo.int/QUARANTINE/EPPO_Study_on_Plants_for_planting.pdf, and Orwig (2002).

²¹ See also the FPS COST Action FP1401 “A global network of nurseries as early warning system against alien tree pests (Global Warning)”, [http://www.cost.eu/COST_Actions/fps/Actions/FP1401].

The impact of invasive alien conifers on hydrology can be enormous, particularly where they replace non-forest vegetation. In South Africa, invasive alien pines were estimated to use 232 million m³ of water per year, about 7 % of water use by all invasive alien plants and about 17 % as much as all commercial forestry (Le Maitre et al. 2000). Run-off in heavily invaded catchments is reduced by 30-70 % (Van Wyk 1987).

Planted forest owners should be aware of those forestry activities that favour the spread of alien tree species outside plantations. For example, coppicing was found to be a driver of the spread by *R. pseudoacacia* and *A. altissima* in South Tyrol, Northern Italy. Radtke et al. (2013) concluded that the currently applied coppice management, which consists of repeated clear cuttings each 20–30 years, favours the spread of both tree species. Thus, they suggest an adaptation of the management system to avoid further spread outside plantations.

Finally, tailored management practices should be followed in the case of planted forests with invasive alien tree for bioenergy production (SRF/SRC), for a careful choice of new planting sites, for favouring biodiversity (Weih 2008; Framstad 2009), protecting hydrology (Christen & Dalgaard 2012), conserving landscape values and for the restoration of the site after the cultivation cycle (Hardcastle 2006; McKay 2011; Neary 2013; Caplat et al. 2014). Development of Forest Management Decision Support Systems for planted forests of alien trees is recommended.

3.2.3 Adopt good practices for harvesting and transport of timber

Install appropriate water and sediment controls and prevent runoff flowing directly into waterways. Keep machinery out of water bodies and riparian margins. Clean and check machinery where the transfer of propagules of invasive alien tree species is an identified risk.

Forest personnel should be trained to recognize and report unusual pests and symptoms of diseased or infested trees, and to carry out practices that reduce the risk of pest and alien species or propagules moving to other locations²². Personnel should wear outer layers of clothing and footwear that are not “seed friendly” (sensu USDA 2012) to minimise the risk of spreading invasive alien species propagules accidentally.

3.2.4 Adopt good practices for habitat restoration

When forest managers wish to remove alien species from forest stands in order to replace them by indigenous species, or when there is wish to remove alien trees from forest stands for other reasons, it is necessary to adopt specific guidelines for the restoration of sites previously occupied by alien trees or by planted forests of alien trees. Restoration objectives can be broadly classified into overarching strategies, such as rehabilitation, reconstruction, reclamation, and replacement (see Stanturf et al. 2014). Native tree species can grow in the understory of planted forests of alien trees. However, not all planted forests of alien tree develop species-rich understories; some remain as alien tree monocultures. Low light intensity below the canopy, distance to seed sources, inhospitability to seed dispersers, poor soil or litter conditions for seed germination or seedling growth, intensive root competition with the planted alien tree species, chemical inhibition and other forms of allelopathy and plant interactions, plantation design, or periodic disturbances by organisms or any external factor are likely causes that require careful consideration (Lugo 1997).

Specific guidelines for restoration of sites previously occupied by planted forests of *Robinia pseudoacacia* have been produced in the Piedmont region of Italy²³. Sturgess & Atkinson (1993) suggested management strategies for the restoration of near-natural sand-dune habitats following the clearfelling of *Pinus* planted forests in Britain, and Brown et al. (2015) proposed approaches for planted forests of alien conifers on ancient woodland sites. Szitár et al. (2014) assessed the recovery of open and closed

²² See also the FAO Guide to implementation of phytosanitary standards in forestry.

²³ <http://www.regione.piemonte.it/foreste/images/files/pubblicazioni/esotiche.pdf>. For Italy see also Maltoni et al. (2012).

grasslands over five years following the removal of planted forests of alien pine species through burning at an inland sand dune system in Hungary. Arévalo and Fernández-Palacios (2005) proposed continuous elimination of *P. radiata* and enrichment with new individuals of *P. canariensis* on Tenerife, Canary Islands (Spain). Hughes (2003) and Moss & Monstadt (2008) propose management guidelines for the restoration of floodplain forests in Europe.

3.3 Early Detection & Rapid Response

3.3.1 *Promote and implement early detection & rapid response programmes*

Early detection and initiation of management can make the difference between being able to employ feasible strategies and facing the necessity of retreating to a more expensive defensive strategy (mitigation, containment, etc.). Proactive measures to reduce the chances of invasive alien tree spread and to deal with problems at an early stage must be incorporated in standard silvicultural practices. Developing alarm lists of possible new invasive alien tree can also enable more rapid reaction (Richardson 2011; Faulkner et al. 2014).

The relatively long initial lag phase between introduction and naturalization/invasion and slow dynamics observed in many invasive alien tree species, in comparison with other plant species, offers opportunities to control the invasive alien species while escaped populations are still small (Finnoff et al. 2007; Dodet & Collet 2012). Any invasive alien tree species detected in the wild or outside cultivation areas should be eradicated, controlled or contained.

Conifer wildings²⁴ of alien species lend themselves to control, as they are relatively easy to detect (most invasions are into grasslands and shrublands), and their direction of spread (downwind), and age when significant seed production begins (usually 10-15 years) is very predictable. There are therefore good opportunities to intercept the spread sequence early in the cycle, and prevent wildings becoming dominant and uncontrollable outside the planted forests (Froude 2011). Natural regeneration of alien conifers is considered desirable in some instances. For example, evolving forest policy in Great Britain requires “lower impact silvicultural systems” on suitable sites that do not require large-scale clearfelling. The transformation of large alien conifer plantations to mixed-aged stands depends to a large extent on natural regeneration (Malcolm et al. 2001). The ecology of natural regeneration, and therefore naturalization and invasion, is thus a highly topical issue (Richardson & Rejmánek 2004). Natural establishment of *E. globulus* plants in Portugal was recently documented by Águas et al. (2014), Catry et al. (2015), Fernandes et al. (2016).

3.3.2 *Establish or join a network of sentinel sites*

Invasive alien trees may be monitored in sentinel site network. Other areas that are likely to act as sources of propagules and sites of entry for new invasions are areas of human habitation where gardens have been established (Alston & Richardson 2006), and experimental plantings, arboreta or botanical gardens containing (invasive) alien tree species. They can be included, as well, in sentinel networks.

²⁴ “Wildings” is the term used (mainly in New Zealand) for the natural regeneration or seedling spread of introduced trees, occurring in locations not managed for forest production. The term is usually applied to members of the family *Pinaceae*, within which most of the major spreading forestry species of concern occur. Most wildings grow close to the parent seed source and are termed fringe spread. Wildings further afield are termed distant spread. They grow from seed often wind-blown from exposed take-off sites and usually occur as scattered outlier trees (Pringle & Willsman 2013). In New Zealand, wilding seedlings are considered vulnerable to grazing for the first 2 years. Mob stocking with sheep will significantly limit their spread, often to the extent that other control requirements are minimal. Cattle grazing is not as effective. Spread can be limited by oversowing and topdressing within a 200-m zone of spread-prone trees. This promotes increased grazing pressure on young wildings and helps the tussock grasslands compete strongly with germinating tree seedlings (“Wilding Prevention” by Nick Ledgard & Lisa Langer, Forest Research Institute, Box 29 237, Fendalton, Christchurch - <http://ecan.govt.nz/advice/your-business/farming/Pages/wilding-trees-preventing-spread.aspx#techniques-prevent-spread>).

Visser et al. (2014) have shown that Google Earth can be a useful tool for establishing a global sentinel site network for alien tree invasions, because imagery is continuously being updated, it is free and low-tech. In addition, the popularity of Google Earth could enable monitoring of this network of sentinel sites as part of a “citizen science” effort (Silvertown 2009). Data sharing via KML files is simple and would allow for easy sharing of locations of sentinel sites. In addition, Google Earth already has the capacity for users to upload photographs (via Panoramio; www.panoramio.com), which would allow for more accurate species identification and verification.

Visser et al. (2014) believe that such a sentinel site network could help to: (1) identify emerging trends in alien tree invasions; (2) provide valuable locality information for particular alien tree species; (3) monitor changes in alien tree species abundance and distribution over time; (4) help ensure legislative compliance of land managers and plantation owners; and (5) track management efforts over time.

In addition to (invasive) alien tree sentinel sites, new technologies such as smartphone application software (apps) are increasingly used to reach a wider audience on the subject of invasive alien species and to involve the public in recording them (Adriaens et al. 2015).

3.4 Outreach

3.4.1 Engage with the public on the risks posed by invasive alien trees, their impacts and on options for management

In addition to land and forest owner, the general public is a very important stakeholder group in national issues of alien trees, from their use in gardening and landscaping to forests and forestry. Public awareness-raising and communication activities play a critical role in informing and educating the public²⁵, thereby allowing them to more effectively participate in decision-making. Public support for control efforts directed at invasive alien trees must be sought through carefully planned, long-term ongoing outreach initiatives involving, among other things, meetings with stakeholders, local village leadership, employment of villagers from areas adjacent to infestations, and the effective use of media outlets.

Forestry has become more complex over the years. This form of land use now benefits a wider stratum of people and environments than ever before, and is subject to a large range of social and environmental demands. As a result, the need for a wide range of professional and managerial skills has increased.

Furthermore, an increasing number of tourists are interested not only in experiencing unique natural and cultural environments and landscapes but also learning more about them. Forest-based tours are an ideal opportunity to share information about different types of forest environments, native and non-native tree species, restoration actions, wildlife and landscapes, how they function and how they came to be. In addition, visitors are also likely to be interested in the lifestyles, cultures and social and political histories of local communities living near forest areas.

²⁵ In Portugal, even though invasive alien species, e.g. acacias, are recognized as a threat to biodiversity by law, the majority of the population is unaware of this problem. Aiming to increase awareness about biological invasions among young students, a workshop on Invasive Plant Species was organized at the Botanical Museum of the University of Coimbra. A total of 170 students from five schools participated in the workshop. Three activities were prepared, focusing on: (1) identification of invasive plants, (2) competition between native and invasive plants and (3) control of invasive plants. A year later, questionnaires were sent to the participants to appraise the effectiveness of the workshop. It revealed that the students know more about invasive plant species than a comparable group of students who did not participate in the workshop. The results clearly showed that practical informal education activities may be effective in raising public awareness. Questionnaires were essential to evaluate the knowledge acquired and retained by the students during the workshop (Marchante et al. 2010; Schreck Reis et al. 2011). See also Andreu et al. (2009); McNeely (ed.) (2001).

3.5 Forward Planning

3.5.1 *Consider developing research activities on invasive alien trees species and becoming involved in collaborative research projects at national and regional levels*

The invasion biology of invasive alien trees is a complex multidisciplinary field and public and private planted forests of invasive alien trees are obvious starting points for field research on topics such as the naturalization, spread, control, management and risks posed by invasive alien trees in collaboration with national or local environment agencies, research centres and appropriate regional or European bodies²⁶.

3.5.2 *Take global change trends into consideration*

Forest management and conservation are expected to be strongly influenced by global change and climate change. Besides forest species, strategies and references for management and conservation will be affected by global change trends²⁷.

Bernier and Schoene (2009) propose three possible approaches for adapting forests to climate change: no intervention, reactive adaptation and planned adaptation²⁸. When applied to planted forests no intervention would mean business as usual, with tree species selection, management targets and practices based on the premise that the planted forest will adapt more or less as it has in the past. Reactive adaptation is action taken after the fact. Planned adaptation, on the other hand, involves redefining forestry goals and practices in advance in view of climate change-related risks and uncertainties. It involves deliberate, anticipatory interventions at different levels and across sectors. Climate change could affect the dynamics of alien²⁹ tree invasions in many interacting ways, for example: (a) by causing modification in the native ecosystems promoting range changes, naturalisation and spread of both native and alien trees

²⁶ E.g., the FPS COST Action FP1403 Non-native tree species for European forests - experiences, risks and opportunities (NNEXT) [http://www.cost.eu/COST_Actions/fps/Actions/FP1403]; the project INVASIVE, Introduced tree species in European forests [<http://www.eficent.efi.int/portal/projects/invasive/>]. INVASIVE is funded by the German Federal Ministry for Food and Agriculture (BMEL) and co-ordinated by the staff at the European Forest Institute (EFI) at the Central European regional office EFICIENT. The partners on the project include FVA Baden-Württemberg, IRSTEA, University of Freiburg, WSL Switzerland and Belgian Biodiversity Platform amongst others. The FA COST Action TD1209 European Information System for Alien Species [http://www.cost.eu/COST_Actions/fa/Actions/TD1209] does not specifically address invasive alien trees, but aims to facilitate enhanced knowledge gathering and sharing through a network of experts, providing support to a European IAS information system which will enable effective and informed decision-making in relation to IAS. An overarching priority will be to identify the needs and formats for alien species information by different user groups and specifically for implementation of EU 2020 Biodiversity Strategy. See also: the FPS COST Action FP1002 Pathway Evaluation and pest Risk Management In Transport (PERMIT) [http://www.cost.eu/COST_Actions/fps/Actions/FP1002].

²⁷ Cf., Jackson et al. (2005); Aitken et al. (2008); Canadell & Raupach (2008); Diaz et al. (2009); Heller & Zavaleta (2009); Thompson et al. (2009); Strassburg et al. (2010); Milad et al. (2013).

²⁸ In the framework of SFM adaptation to climate change is the “adjustment in natural or human systems in response to actual or expected climatic stimuli or their effects, which moderates harm or exploits beneficial opportunities. Various types of adaptation can be distinguished, including anticipatory and reactive adaptation, private and public adaptation, and autonomous and planned adaptation”. Ecosystem-based adaptation to climate change is an adaptation strategy that includes ecosystem management, conservation and the restoration of ecosystems to provide services that help people adapt to adverse impacts of climate change. Cf. also McCarthy et al. (2001); Locatelli et al. (2008); CBD Secretariat (2009). Connecting biodiversity and climate change mitigation and adaptation: report of the Second Ad Hoc Technical Expert Group on Biodiversity and Climate Change. Technical Series No.41. CBD Secretariat, Montreal, Canada.

²⁹ In accordance with the Recommendation No. 142 (2009) of the Standing Committee, adopted on 26 November 2009, interpreting the CBD definition of invasive alien species to take into account climate change, the term “alien tree species” for the purpose of this Code of Conduct does not include native tree species naturally extending their range in response to climate change.

(e.g., Iverson et al. 2008; McKenney et al. 2011); (b) by favouring individual traits of particular alien trees (e.g. Capdevila-Argüelles & Zilletti 2008; Kawaletz et al. 2013; Castro-Díez et al. 2014); and (c) by modifying introduction pathways and promoting a larger use of certain alien trees (Courbet et al. 2012; Lindenmayer et al. 2012) including a process of re-thinking the importance of the “always choosing native species” principle (UK Forestry Commission³⁰). Managed relocation³¹ has been proposed as a means to maintain forest productivity, health, and ecosystem services under rapid climate change (e.g., Gray et al. 2011; Kreyling et al. 2011; Pedlar et al. 2012).

Finally, it is important to incorporate climate change into risk assessment and risk analysis models for an anticipatory evaluation of scenarios for invasiveness of alien trees. Risk maps³² that incorporate the effects of climate change should help land managers and forest stakeholders with longer-term planning activities. Management plans of nature reserves should incorporate changes to invasion risk driven by global warming more explicitly.

4. BACKGROUND INFORMATION: A KNOWLEDGE BASIS FOR THE CODE OF CONDUCT

4.1 Benefits arising from planted forests and planted forests of alien trees

Planted forests are forests predominantly composed of trees established through planting and/or deliberate seeding (FAO 2012)³³. Planted forest of introduced species (PFIS) is a subcategory of planted forest, where the planted/seeded trees are predominantly of introduced species (FAO 2012). In 2010, the total area of planted forests was estimated to be 264 million ha (about 7 % of the total global forest area; FAO 2010a), and this increased to an estimated 277.9 million ha in 2015 (FAO 2015a, 2015b; Payn et al. 2015)³⁴. Although there are marked differences between and within regions, it has been estimated that between 18% and 19% of planted forests comprise alien tree species (Payn et al. 2015; FAO 2015a, 2015b).

Planted forests designed to provide multiple ecosystem services can reduce pressure on natural forests, and can even restore some ecological services provided by natural forests. They can also play a key role in the fight against global warming, through carbon sequestration. Paquette & Messier (2010) reviewed the economic, social, and environmental services that plantations can provide, and make a plea for the implementation of well-conceived, diverse, multi-purpose plantations as a way to conserve forest

³⁰ [http://www.forestry.gov.uk/pdf/eng-trees-and-climate-change.pdf/\\$FILE/eng-trees-and-climate-change.pdf](http://www.forestry.gov.uk/pdf/eng-trees-and-climate-change.pdf/$FILE/eng-trees-and-climate-change.pdf). See also Forestry Commission (2011); NOMADES, *NOuvelles Méthodes d'Acclimatation Des Essences forestières* (<http://www.reseau-aforce.fr/nomades-437950.html>) - http://www.reseau-aforce.fr/data/info/497950-NOMADES_Fascicule1_Bilan_introduction_vdef_fev15.pdf); REINFFORCE, REsource INFrastructures for monitoring, adapting and protecting european atlantic FORests under Changing climatE (http://www.iefc.net/?affiche_page=projet_REINFFORCE&langue=en).

³¹ Managed relocation or assisted migration has been proposed as an approach to mitigate climate change impacts on biodiversity by intentionally moving species to climatically suitable locations outside their natural range (Richardson et al. 2009).

³² Pest risk maps are powerful visual communication tools to describe where invasive alien species might arrive, establish, spread, or cause harmful impacts. These maps inform strategic and tactical pest management decisions, such as potential restrictions on international trade or the design of pest surveys and domestic quarantines. Diverse methods are available to create pest risk maps, and can potentially yield different depictions of risk for the same species (Venette et al. 2010).

³³ In the context of the 2015 FAO Global Forest Resources Assessment (FRA), predominantly means that the planted/seeded trees are expected to constitute more than 50 percent of the growing stock at maturity. In addition, planted forest: (a) includes coppice from trees that were originally planted or seeded, (b) includes rubberwood, cork oak and Christmas tree plantations, (c) excludes self-sown trees of introduced species.

³⁴ Planted forest area has increased by over 110 million ha since 1990 and accounts for 7 percent of the world's forest area. The average annual rate of increase between 1990 and 2000 was 3.6 million ha. The rate peaked at 5.2 million ha per year for the period 2000 to 2010 and slowed to 3.1 million ha (2010–2015) per year, as planting decreased in East Asia, Europe, North America, South and Southeast Asia (FAO 2015 b).

biodiversity and ecosystem functions. Well-designed, multi-purposed planted forests can help mitigate climate change³⁵ through direct carbon sequestration or by avoiding deforestation, while simultaneously protecting remaining natural forests through increased productivity.

Standards, guidelines, criteria and indicators for sustainable forest management (SFM) have been developed over the past few decades by intergovernmental processes, international organizations³⁶, certification schemes (e.g. FSC, PEFC) (Masiero et al. 2015) and national governments (Cf. Section 4.6.6 in this Code of Conduct). These apply to all forests, including planted forests, and have resulted in forestry being recognised as an essentially sustainable land-use and essential to combatting climate change by storing carbon and preventing deforestation. Activity was increased considerably after the Statement of Principles for the Sustainable Management of Forests was adopted in 1992 at the Earth Summit in Rio in response to global concerns about deforestation and the unsustainable exploitation of natural forests (Stupak et al. 2011). At European level, the 46 signatories of the Ministerial Conference on the Protection of Forests in Europe agreed a definition of sustainable forest management in a Ministerial Process dating from 1990 and have developed and refined a set of criteria and indicators. These are regularly updated and adapted to new challenges³⁷. The International Tropical Timber Organization ITTO has developed and is revising criteria and indicators for sustainable forest management since the early 1990's³⁸.

Recognizing the economic, social, cultural and environmental importance of planted forests, governments and other stakeholders asked FAO to prepare, together with collaborating partners, a set of guiding principles in support of the policy, legal, regulatory and technical enabling conditions for planted forest management (FAO 2006b).

Since 1980, the Food and Agriculture Organisation of the United Nations (FAO) through its Forest Resources Assessments (FRA), has been collecting data on forest areas for two main categories of forests: natural forests and forest plantations³⁹. In 2005, the FRA introduced two additional forest categories: modified natural forests and semi-natural forests (Evans 2009a), which resulted in five major forest categories based on the degree of human intervention and the silvicultural methods of forest regeneration. These include (1) primary forest; (2) modified natural forest; (3) semi-natural forest, comprising natural and planted regeneration (SNPF); (4) plantations comprising productive and protective plantations; and (5) trees outside forests (Payn et al. 2015). Productive and protective plantations, together with SNPFs,

³⁵ E.g., for France see Blondet M. (2015), *Conflicting engagements on climate change adaptation in French private forest: an anthropological perspective*. Document de travail du LEF n°2015-03 (LEF, AgroParisTech/INRA, Nancy, France). The issue of climate change is progressively entering the field of forest management in France and Europe. It poses significant questions to forest managers since forest management is made on a very long-time scale. Decisions taken today will impact forest for many years and climate change may threaten these long-term investments. According to scientists, beech forest is particularly sensitive to drought and may disappear in the coming years due to global warming. Beech is also one of the protected species in the Annexes of the Habitat Directive. To face and bring answers to this issue of the future of beech forest before this change in climate conditions various actors from the forest sector, the conservationist organisations and the policy-making sphere are engaging at the national level in France. Yet they carry different views of the issue. In France, and within that frame, a debate recently started around the possible use of planted forest of alien trees to control the impacts of climate change in forest. The idea basically is to seek for more resistant tree species in order to replace the weakest ones such as European beech. These stronger species are often not indigenous species, which may be subject to controversy among actors from the forest sector, the conservationist organisations and the policy-making sphere.

³⁶ Cf., *New Generation Plantations (NGP)*, 2014. *New generation plantations: review 2014* (<http://newgenerationplantations.org/multimedia/file/12b486cb-ea24-11e3-9f9e-005056986313>).

³⁷ Forest Europe http://www.foresteuropa.org/sfm_criteria/criteria

³⁸ ITTO http://www.itto.int/sustainable_forest_management/

³⁹ In the FAO FRA 2000 (FAO 2001) "forest plantations" were defined as those forest stands established by planting or/and seeding in the process of afforestation or reforestation. They comprised either native or non-native species which met a minimum area requirement of 0.5 ha; tree crown cover of at least 10 % of the land cover; and total height of adult trees above 5 m.

constituted the subgroup “planted forests”⁴⁰ (FAO 2010a). In the Global Forest Resources Assessment 2010 (FAO 2010a) the concept of “planted forests” was defined more broadly than the concept of forest plantations as used in previous global assessments. This change was made to capture all planted forests and is in line with the recommendations of the Global Planted Forests Thematic Study 2005 (FAO 2006a) and recent efforts to develop guidelines and best practices for the establishment and management of planted forests. The FRA 2015 definition (FAO 2012) refined this to: forest predominantly composed of trees established through planting and/or deliberate seeding, where the planted/seeded trees are expected to constitute more than 50 % of the growing stock at maturity. They include coppice from trees that were originally planted or seeded and rubber wood, cork oak and Christmas tree plantations (Payn et al. 2015).

East Asia, Europe and North America hold the greatest area of planted forests, together accounting for about 75 % of global planted forest area, followed by North America and Southern and Southeast Asia (FAO 2010a; Payn et al. 2015). In East Asia planted forests make up 35 % of the total forest area; most of these are found in China. The second largest area of planted forests is found in Europe, although the share of planted forests here is close to the world average. However, if the Russian Federation with its vast area of natural forest is excluded from Europe, the share of planted forests in Europe increases to 27 %, the second highest proportion in the world. North America has the third largest area of planted forests with 5.5 % of the total forest area occurring in this subregion. Subregions reporting the smallest area of planted forests are the African subregions, the Caribbean, Central America and Western and Central Asia. In most subregions, the majority of the planted forests are found in just a few countries. For instance, in northern Africa 75 % of the planted forest area is located in Sudan, in East Asia, 86 % is found in China. Some arid-zone countries (Cape Verde, Egypt, Kuwait, Libyan Arab Jamahiriya, Oman, United Arab Emirates) and the Netherlands report that all their forests have been established through planting or deliberate seeding (FAO 2010a).

Between 2000 and 2010, the area of planted forest increased by about 5 million ha per year (FAO 2010a), with a further increase in the period 2010-2015 (Payn et al. 2015; FAO 2015b). Most of this was established through afforestation, particularly in China.

4.2 Alien tree species in Planted Forests and for other uses: historical and recent pathways of introduction

4.2.1 General aspects

As stated in Section 1, in accordance with the CBD and the Reg. (EU) No. 1143/20014, the term alien tree is used throughout this Code of Conduct. It has the same meaning as exotic, introduced and non-native. In accordance with the CBD definition, the term alien tree has exclusively a biogeographical meaning, i.e. it refers to a tree species, subspecies or lower taxon (including provenance), introduced by man outside its natural past or present distribution.

After the United Nations Conference on Environment and Development (UNCED) in 1992, known as the Rio Summit, an enhanced understanding of sustainable forest management (SFM) has entered the stage of forest policy worldwide (Wolfslehner et al. 2005; Cf. Section 4.6.6 in this Code). Key outcomes of the conference were the “Forest Principles” and the Chapter 11 of Agenda 21 of the conference’s action plan (UNCED, 1992a, 1992b, 1992c) which detailed the international commitments to the sustainable development of forests, and called for the formulation of scientifically sound criteria and indicators (C&I) for sustainable forest management (Baycheva-Mergera & Wolfslehner 2016).

⁴⁰ In Annex 2, “Terms and definitions used in FRA 2010” page 212, *planted forest* is defined as follows: Forest predominantly composed of trees established through planting and/or deliberate seeding; the sub-category *planted forest of introduced species* as planted forest, where the planted/seeded trees are predominantly of introduced species.

However, in addition to the use in planted forests, alien tree species have been and are introduced and used for various and multiple reasons (i.e. introduction pathways⁴¹), such as gardening, protective functions, arboreta, erosion protection and for increasing the forest area through afforestation of abandoned or derelict land.

In the 2010 FAO Global Forest Resources Assessments (FAO 2010a) countries reported on the use of introduced⁴² tree species in the establishment of planted forests of introduced species. Globally, of the 233 countries and areas included, only 117⁴³ countries reported on the use of introduced species, while the remaining 116 countries and areas did not report on the use of introduced species.

At the global level, introduced tree species grow on about a quarter of the planted forest area of the countries for which data were reported (FAO 2010a). Payn et al. (2015), using FRA 2015 datasets (FAO 2015a, 2015b), estimated that only between 18% and 19% of the planted forests comprise introduced tree species. There are marked differences between and within regions. South America, Oceania, and East and Southern Africa are the regions dominated by plantings of introduced tree species, with large percentages of plantings comprising introduced tree species. North America, West and Central Asia, and Europe are at the other end of the spectrum with 1%, 3% and 8% of the area planted in introduced tree species Payn et al. (2015).

In eastern and southern Africa, most planted forests consist of introduced tree species in the genera *Eucalyptus*, *Pinus*, *Hevea*, *Acacia* and *Tectona*, chosen for their ability to grow in many environmental conditions and to rapidly produce wood or other economic products (e.g., gum Arabic, rubber) (FAO 2010a).

Planted forest species in Oceania and in South America (Argentina, Bolivia, Brazil, Chile, Ecuador and Uruguay) also comprise introduced tree species. Oceania has a long history of planted forest management due to historic wood supply deficits and offers excellent growing conditions for a number of fast growing species, among them *Eucalyptus spp.*, *Pinus radiata* (Monterey Pine), *Pseudotsuga menziesii* (Douglas-fir) and *Cupressus spp.*⁴⁴. South America is encouraging the use of intensively managed short-rotation introduced tree species such as *Eucalyptus spp.*, *Pinus radiata*, *P. taeda*, *P. elliottii* and *Tectona grandis*.

⁴¹ The term **pathways** is used in accordance to the CBD definition, i.e. UNEP/CBD/SBSTTA/18/9/Add.1 of 26 June 2014, Pathways of Introduction of Invasive Species, their Prioritization and Management (<https://www.cbd.int/doc/meetings/sbstta/sbstta-18/official/sbstta-18-09-add1-en.pdf>). The Guiding Principles for the Prevention, Introduction and Mitigation of Impacts of Alien Species that threaten Ecosystems, Habitats and Species (the Guiding Principles) annexed to decision VI/23, provide all Governments and organizations with guidance for developing effective strategies to minimize the spread and impact of invasive alien species. In particular, the Guiding Principles highlight the importance of identifying pathways of introduction of invasive species in order to minimize such introductions, and call to assess the risks associated with such pathways. Furthermore, Aichi Biodiversity Target 9 specifies: “By 2020, invasive alien species and pathways are identified and prioritized, priority species are controlled or eradicated, and measures are in place to manage pathways to prevent their introduction and establishment”.

⁴² The 2010 FAO Global Forest Resources Assessments uses the term “introduced”. The definition of “introduced species” is as follows: a species, subspecies or lower taxon, occurring outside its natural range (past or present) and dispersal potential (i.e. outside the range it occupies naturally or could occupy without direct or indirect introduction or care by humans). The FAO 2015 Global Forest Resources Assessment (FRA), defines an introduced species as a species, subspecies or lower taxon, occurring outside its natural range (past or present) and dispersal potential, i.e. outside the range it occupies naturally or could occupy without direct or indirect introduction or care by humans, FAO (2012). In FRA 2015 the term introduced is considered equivalent to non-native.

⁴³ Seventeen countries reported that they have not used introduced species in the establishment of planted forests (FAO 2010).

⁴⁴ *Cupressus macrocarpa*, *C. lusitanica*, and the closely related *C. benthamii*.

In East Asia, China uses introduced tree species on 28 % of the planted forest area while Japan did not report the proportion of introduced tree species (FAO 2010a, 2015a). In South and Southeast Asia, a number of countries with a significant area of planted forest did not report on the use of introduced tree species (Indonesia, Malaysia and Vietnam⁴⁵). In Southeast Asia, plantations are established more for non-timber crops than timber, particularly coconuts, rubber, and oil palm (Corlett 2005) but there is a growing interest for Acacia and Eucalyptus plantations (Harwood & Nambiar 2014).

In western and central Asian countries (e.g., Turkey⁴⁶ - 59,000 ha, FAO 2015a) the use of introduced tree species is very low, while other countries in this subregion did not report on introduced tree species. In the temperate and boreal regions of Europe and North America and in the arid zone countries of northern Africa introduced tree species are only used to a minor extent.

Some parts of Europe lack highly productive native tree species with timber or growth characteristics suited to planted forests, and foresters rely largely upon non-native tree species. These alien tree species can be established easily on certain sites, have better growth rates than native species, broader physiological adaptability with regard to site conditions including drought tolerance (Savill et al. 1997). The area dominated by introduced tree species covers about 9 million ha or 4 % of the total forest area (without the Russian Federation). In the Russian Federation, less than 100,000 ha of its vast forest area was reported as planted forest with non-native trees, thus being negligible (66,000 in 2015, FAO 2015 a). In Denmark, Iceland and Italy⁴⁷, introduced tree species are reported to occur also on other wooded land (Forest Europe, UNECE and FAO 2011).

The most important alien tree species used in Europe for timber production include *Pseudotsuga menziesii*, *Picea sitchensis*, *Pinus contorta*, *Populus* hybrids and clones, and a number of *Eucalyptus* spp. The relative absence of pests and specialized grazing or defoliating insects from aliens allows the trees to grow much faster than native species until pests catch up with their hosts, especially if unaccompanied by their natural enemies (Savill et al. 1997). However, when plantation alien trees are reunited with their coevolved pests, which may be introduced accidentally, or when they encounter novel pests to which they have no resistance, substantial damage or loss can ensue. The longer these non-native trees are planted in an area, the more threatened they become by native pests (Wingfield et al. 2015).

In Sweden, Elfving & Norgren (1993) have demonstrated that *Pinus contorta*⁴⁸ (lodgepole pine) can grow 32 % faster in terms of stemwood volume than the native *P. sylvestris* (Scots pine), because the former allocates more resources to the growth of stems and fine roots rather than larger roots compared to the latter (Savill et al. 1997). Other reasons for the superior growth of lodgepole pine under boreal conditions may be an earlier start of growth in spring and a lower required heat sum to start shoot elongation compared to Scots pine (Fedorkov 2010; Backlund & Bergsten 2012). Despite the apparent growth, and hence economic, benefits of alien tree species, fears of eventual pest and disease outbreaks led to legislation in 1979 (Swedish Forestry Act) and in 1992 aimed at limiting the use of *P. contorta* until the potential risks are better understood (Savill et al. 1997).

⁴⁵ According to IUFRO (Scientific Summary No 120, related to IUFRO News 4, 2014), the most significant areas of plantations of Australian Acacias are in SE Asia where *A. mangium*, its hybrid with *A. auriculiformis* and *A. crassiparpa* are the main taxa. In Vietnam over 1M ha of Acacia plantations supply a burgeoning furniture manufacturing industry as well as the export woodchip market.

⁴⁶ The Strategic Plan of the General Directorate of Forestry (2013-2017) is a commitment to industrial plantations with fast growing species. It is planned that industrial plantations are established in a total area of 15,000 ha by the end of 2014 (Deniz & Yildirim 2014).

⁴⁷ Between the 1920 and the 1939, 450 experimental plots for 124 non-native tree species were established in Italy with the purpose of comparing their productivity performances (Pavari & De Philippis 1941; Nocentini 2010).

⁴⁸ *Pinus contorta* Dougl. var. *latifolia* Engelm.

Pseudotsuga menziesii was introduced to Europe from North America more than 150 years ago⁴⁹, and is now the most economically-important alien tree species in European planted forests (Schmid et al. 2014). It was introduced to Sweden in the 1920s, and its planted forests are currently estimated to occupy approximately 500 ha, primarily on large estates in southern Sweden (Felton et al. 2013). Broncano et al. (2005) described the naturalization by *Pseudotsuga menziesii* in Montseny Natural Park (Catalonia, NE Spain – a declared a UNESCO Biosphere Reserve in 1978). Establishment of seedlings started 15 years after plantings. Essl (2005) reports the naturalization of *Pseudotsuga menziesii* in lowland northeastern Austria, and there are naturalized occurrences in most other Central European countries (Schmid et al. 2014)⁵⁰.

Northern red oak (*Quercus rubra*) is an economically important, moderately shade-tolerant tree species native to eastern North America (Sander 1990). While this species is failing to regenerate in many locations in its native range, red oak has regenerated readily in Central European forests since its introduction in the mid-18th century (Kuehne et al. 2014). The ability of non-native red oak to perform equally well to native shade-tolerant species under a variety of light conditions could contribute to the consistent success of red oak regeneration in Europe (Riepšas & Straigītē 2008; Kuehne et al. 2014; Woziwoda et al. 2014).

Robinia pseudoacacia has been widely used for various purposes such as ornamentation, timber (wood quality comparable to hardwood tropical trees), firewood, re-vegetation of dry land, soil stabilisation, fertilisation of poor soils and providing nectar for honey production (EEA 2008). *Robinia pseudoacacia* stands occupy 20 % (about 400,000 ha) of the Hungarian forest area (Rédei 2002; Rédei et al. 2011b), and it has been documented that it is a pioneer species⁵¹. Having only one native species, downy birch (*Betula pubescens*), and a small forest area, Icelandic forests have a high proportion of introduced tree species due to afforestation efforts.

Sitka spruce (*Picea sitchensis*) from North America is the most common alien tree in Great Britain and Ireland (Peterken et al. 1992; Quine & Humphrey 2010; Peterken 2001). Britain has a very limited native tree flora, but now has a great variety of introduced trees (Peterken 2001). The most commonly used alien coniferous trees are *Abies alba*, *Larix decidua*, *Picea abies*, *Picea sitchensis*, *Pinus contorta*, *Pseudotsuga menziesii*, *Tsuga heterophylla*. Introduced trees have formed hybrids in Britain, both with native trees and amongst themselves. *Quercus* × *turneri* (syn. *Quercus* × *hispanica*) is a semi-evergreen oak formed in the late 18th century by *Q. ilex* × *Q. robur*.

Ailanthus altissima, mainly used as an ornamental or for roadside plantings, is considered one of the most widespread invasive alien plant species in Europe and North America (Sladonja et al. 2015). *Acer negundo* and *Prunus serotina*⁵² are both ranked third and are reported as invasive alien trees in several European countries (Forest Europe, UNECE and FAO 2011).

⁴⁹ David Douglas introduced *Pseudotsuga menziesii* to Great Britain in 1827 (Gellini & Grossoni 1996).

⁵⁰ Cf.: Tschopp, T., Holderegger, R., Bollmann, K., 2014: Auswirkungen der Douglasie auf die Waldbiodiversität : Eine Literaturübersicht. WSL Berichte, Heft 20, 2014 (55 S.); Tschopp, T., Holderegger, R., Bollmann, K., 2015: Auswirkungen der Douglasie auf die Waldbiodiversität. Schweiz. Z. Forstwesen 166 (2015) 1: 9-15: Vor, T., Spellmann, H., Bolte, A., Ammer, C. (Hrsg.), 2015: Potenziale und Risiken eingeführter Baumarten. Baumartenportraits mit naturschutzfachlicher Bewertung. Deutscher Verband forstl. Forschungsanstalten, 2015 (233 S.): <http://www.dfwr.de/aktuelles/> -> Rubrik "Aktuelles aus der Forstwirtschaft", Mitteilung vom 23. Februar 2015, Dokument Nr 5.

⁵¹ Pure or mixed stands of *Robinia pseudoacacia* now cover some 200,000 ha in France, 250,000 ha in Romania and 230,000 ha in Italy (Sitzia 2014).

⁵² *Prunus serotina* (syn. *Padus serotina*), a forest tree of North American origin (Mexico-Guatemala to south-east Canada), was introduced to central Europe and planted for various purposes. The first record of the species in Europe dates back to 1623 when the tree was planted for ornamental purposes near Paris (Starfinger 1997), in England in 1629 and in Germany in 1685 (Starfinger & Kowarik 2003), where it was particularly appreciated as an ornamental (Petitpierre et al. 2009). Between 1900 and 1930, black cherry was planted for multiple uses such as wind and firebreaks, to improve soils under coniferous plantations, or for shelter (Pairen et al. 2010; Vanhellemont et al.

A shift in forest management to increase the share of native tree species has led to a steady decline of introduced tree species (e.g., in the Netherlands). Countries with a very low share, i.e. below 0.5 %, of introduced tree species or no introduced tree species are Lithuania, Finland, Estonia, Serbia, Latvia, Belarus, Liechtenstein and Georgia (Forest Europe, UNECE and FAO 2011, 2015a).

4.2.2 Alien conifers

Many alien conifers are very widely used in planted forests, and as amenity and ornamental plants (Richardson & Rejmánek 2004). *Pinus radiata*, from a tiny native range in California and a few islands, has been planted over huge areas in alien plantations, mostly in the southern hemisphere, especially New Zealand, Chile, Australia and South Africa. A total number of 38 Conifers are listed in the global database of invasive trees and shrubs (Rejmánek & Richardson 2013), with 15 of them indicated as invasive also in Europe⁵³.

In general, conifer taxa from Europe and North America have been more widely planted well outside their natural ranges than those from other regions, notably Asia (Richardson & Rejmánek 2004 and references cited therein). An example is *Picea asperata* and its close relative *P. abies*. *P. asperata* (from China) has enjoyed trivial planting and dissemination outside its range compared to the European *P. abies*. Several *Pinus* species are among the most widespread and influential of all invasive alien trees, especially in the southern hemisphere (Richardson & Rejmánek 2004).

There are 56 non-native coniferous species recorded in Denmark. Seven of these are regarded as invasive and they are already on the Black list (see Annex 6.2). All were introduced intentionally for forestry and horticulture (Madsen et al. 2014).

Norwegian forestry has mainly used two native coniferous tree species, *Picea abies* and *Pinus sylvestris*, although attempts have been made to plant alien tree species, some of which are in current use (Felton et al. 2013). All alien tree species that have been planted have produced seed, and many have spread to a lesser or greater degree outside plantations. Only 4–5 of these alien species are considered as problem species to any degree, yet the fact that they can alter the environment rather dramatically where they become established means that they can locally, and perhaps also regionally, have marked impacts on biodiversity (Gederaas et al. 2012). The introduced species include *Picea sitchensis*, *Tsuga heterophylla*, *Pinus contorta*, *Larix decidua* and *Pinus mugo*. One of the species which is considered to have a severe impact (SE) is *Picea sitchensis*, and it is included in the Norwegian Black-List (see Annex 6.2).

2010). It was first considered a valuable timber tree by European foresters, then a useful non-timber species in forestry, then a forest pest, a controllable weed, an invasive alien tree and, eventually, a species we have to live with. All these perceived qualities served as motives for action by humans often without seeking scientific evidence for them: millions of specimens of *P. serotina* were planted. Later millions of Euros were spent in attempts at control (e.g., Van den Meersschaut & Lust 1997). The overall loss to the German economy through yield reduction and control costs was estimated at 25 million of Euros per year. A similar figure was estimated for the Netherlands (Starfinger 2010). The species, and its changing perception through time, may be an example of the need for science-based assessments as a basis for developing policies concerning the use of non-native tree species in planted forests (Starfinger et al. 2003). In Italy, the Lombardy Region warns about the risks posed by *Prunus serotina* in planted forests (<http://www.arpalombardia.it/biodiversita/piante-viaggiatrici/landing-schede/prunus-serotina.html>). According to the most recent studies, *Prunus serotina* grows spontaneously in many European countries from France to Poland and Romania as well as from Denmark to Italy. It is spreading throughout temperate forests in north-western Europe, especially on well-drained poor soils. This tree is now considered as one of the 100 most invasive alien species in Europe (DAISIE, 2009). *Prunus serotina* competes for resources with native plant species, especially during forest regeneration and under high herbivore pressure. Invaded stands have higher levels of phosphorus, a shallower litter layer, and lower pH values than non-invaded stands. Shading out light-demanding species, especially seedlings of other tree species, it can impede natural regeneration of native tree species and induces a decrease in species richness, mainly in disturbed stands. Fruits are produced in high quantities and are well dispersed, mainly by birds. More than 50% of the seeds are dispersed at higher distances than 50 m; the average dispersal distance being estimated to 257 m (Pairon et al. 2010 and references cited therein).

⁵³ See also: Carrillo-Gavilán and Vilà (2010).

The North American tree *Pinus contorta* var. *latifolia* was experimentally introduced to Sweden as early as the 1920s, and has been used in Swedish forestry on a large scale since the 1970s. Plantations of this species now cover 475,000 ha (with at least 65% Lodgepole pine), mainly in the northern part of the country (Engelmark et al. 2001; SLU 2010).

In Iceland, due to the lack of native trees suitable for planted forests, alien tree species (and conifers in particular) are economically important. Numerous conifers were introduced and are in use in Icelandic forestry, but *Pinus contorta* and *Picea sitchensis* have already become naturalised and started to spread outside cultivation (Wasowicz et al. 2013).

4.2.3 Alien eucalypts

Over 800 species of eucalypts (the genera *Angophora*, *Corymbia* and *Eucalyptus*) are native to Australia and a few Pacific islands. These genera include some of the most important solid timber and paper pulp alien forestry trees in the world. Besides pines, eucalypts are the most commonly and widely cultivated alien trees. Over 70 species are naturalised alien (reproduce and maintain their populations) outside their native ranges. However, given the extent of their cultivation, eucalypts are markedly less naturalized and less invasive than many other widely cultivated alien trees and shrubs. Reasons for this relatively low invasiveness are not completely understood (Rejmánek & Richardson 2011; Rejmánek and Richardson 2013; Águas et al. 2014; Catry et al. 2015; Lorentz & Minogue 2015).

Eucalypts have been planted for forestry over large areas in Spain and Portugal, and to a more limited extent in Italy and Turkey. In Spain, slightly more than 3.5 % of the total forestry area (Anuario de estadística forestal 2011) and in Portugal about 812,000 ha (*Inventario Florestal Nacional 5, 2005-06, 2013*) are covered by *Eucalyptus* species (Forest Europe, UNECE and FAO 2011). Two species have been recorded as naturalised alien in Europe (*E. camaldulensis* and *E. globulus*, e.g. Catry et al. 2015).

4.2.4 Alien acacias

Like pines and eucalypts, many acacias (a polyphyletic group comprising more than 1,350 species, according to Maslin et al. 2003), and especially Australian acacias, have been widely planted outside their natural ranges for centuries (Richardson et al. 2011; Kull et al. 2011). Alien acacias besides being commercially important crops, play diverse roles in the lives and livelihoods of rural communities around the world. Landscapes in many parts of the world are dominated by planted or self-sown stands of Australian acacias. Some species are crops of major commercial importance and many others have considerable value for a wide range of purposes. Some Australian acacias are among the most widespread and damaging of all invasive alien plants (Richardson & Rejmánek 2011). Others are considered only moderately invasive, and yet others are not known to invade, although some of the last-mentioned are recent introductions.

A. melanoxylon was introduced in Portugal as an ornamental in the mid-nineteenth century and its expansion occurred in the first half of the twentieth century through national reforestation programmes, in which the afforestation projects of coastal dunes included aliens such as *Acacia*, *Casuarina* and *Eucalyptus* (Goes 1991; Leite et al. 1999; Knapic et al. 2006). In Spain, *A. melanoxylon* is currently widely naturalized in Galicia, northern Spain, in areas below 500 m altitude, usually in low sloping lands (Sanz-Elorza et al, 2004). *Acacia dealbata* is on the Black List for Spanish mainland, and *A. farnesiana* and *A. salicina* for Canary Islands⁵⁴. *Acacia dealbata* has been described as a serious environmental problem in Northwest Spain, where its expansion is assumed to reduce populations of native species and threaten local plant biodiversity (Lorenzo et al. 2010, 2011).

⁵⁴ *Real Decreto 630/2013, de 2 de agosto, por el que se regula el Catálogo español de especies exóticas invasoras* (Act 630/2013, 2nd August, that regulates Spanish Catalogue on Invasive Alien Species - <http://www.boe.es/buscar/act.php?id=BOE-A-2013-8565>). Cf. also García-de-Lomas & Vilà (2015).

A total of 33 species in the genus *Acacia* are listed in the global database of invasive alien trees and shrubs (Rejmánek & Richardson 2013); 9 species are considered invasive alien in some European countries (8 species from Australia and one species from Africa)⁵⁵.

4.2.5 *Alien poplars and willows*

An estimated 70 countries grow poplars and willows in mixtures with other natural forest species, in planted forests and as individual trees in the landscape (including agroforestry systems). Country reports to the International Poplar Commission (IPC⁵⁶) indicate that poplars and willows account for more than 95 million ha of natural (82 million ha) and planted forests and agroforestry production systems (13 million ha) globally. The Russian Federation, Canada and the United States have the largest reported areas of naturally occurring poplar and willows, while China, India and Pakistan have the largest planted areas (FAO IPC Website 2014⁵⁷).

Poplars and willows are multi-purpose species and form an important component of forestry and agricultural production systems worldwide, often owned by small-scale farmers. They provide a long list of wood and fibre products (sawn lumber, veneer, plywood, pulp and paper, packing crates, pallets, poles, furniture and small handicraft), non-wood products (animal fodder), environmental services (rehabilitation of degraded lands, forest landscape restoration, climate change mitigation) and are grown increasingly in bio-energy plantations for the production of biofuels. These attributes make poplars and willows ideally suited for supporting rural livelihoods, enhancing food security, alleviating poverty and contributing to sustainable land-use and rural development (FAO IPC Website 2014).

Transgenic poplars have been used in numerous regulated field trials in the USA (Strauss et al. 2004), and are currently being commercially cultivated only in China (Sedjo 2005). There is a large potential for additional transgenic applications because the poplar genome has been sequenced, many genotypes are amenable to genetic transformation, and transformation appears capable of improving its high value for bioremediation as well as a number of other traits (Boerjan 2005; Di Fazio et al. 2012). Poplars are dioecious and wind-pollinated, and produce abundant, small seeds with cotton-like appendages that facilitate long-distance dispersal by wind and water. Finally, wild relatives are often interfertile with cultivated clones, and extensive wild populations commonly occur in the vicinity of commercial plantations (Di Fazio et al. 2012). There are substantial concerns about the spread of transgenic plants into wild and feral plant communities. These concerns are heightened for perennial species such as trees that have undergone little domestication and that provide extensive ecological services (James et al. 1998; Hoenicka & Fladung 2006; Di Fazio et al. 2012).

North American feltleaf willow (*Salix alaxensis*), recently introduced as a forestry species in Iceland, has become naturalised and started to spread very effectively. It seems that the spread of this species will be further facilitated by climate change (Wasowicz et al. 2013).

Five *Populus* and 14 *Salix* taxa are listed in the global database of invasive trees and shrubs (Rejmánek & Richardson 2013); three *Populus* (*P. alba*, *P. × canadensis*, *P. × canescens*) and two *Salix* taxa (*S. daphnoides*, *S. fragilis*) are listed as being invasive to and in Europe (i.e., although native in some parts of Europe).

⁵⁵ See also Pasta et al. (2012) for the Mediterranean region. Vietnam has over 400,000 ha of *Acacia* plantations, including over 220,000 ha of clonal *Acacia* hybrid (*Acacia mangium* × *Acacia auriculiformis*). *Acacia* hybrid has been planted extensively in the southern provinces of Vietnam, and is becoming one of the main species for industrial plantations (Sein & Mitlöhner 2011).

⁵⁶ The International Poplar Commission (IPC) is one of the oldest statutory bodies within the framework of the Food and Agriculture Organization of the United Nations (FAO). It was founded in 1947 by 9 European countries in the aftermath of WWII destructions, when poplar and willow culture was considered a priority to supporting reconstruction of rural and industrial economies (<http://www.fao.org/forestry/ipc/en/>).

⁵⁷ <http://www.fao.org/forestry/ipc/69994/en/>

4.2.6 *Alien trees and plantations on disturbed land*

Numerous industrial processes disturb land of which the principal ones are mining, extraction of sand, gravel and clay, rock and limestone quarries, deposition of waste products including landfill sites, road and railway construction. The greatest amount of dereliction occurs in industrial countries. The problem arises principally because the substrate to be reclaimed is almost always derived from mining or earth moving, and it is largely undeveloped subsoil or rock or it is polluted. The nature of reclaimed sites necessitates the use of species which are tolerant of exposure and undemanding nutritionally, characteristics often associated with pioneer species including alien trees (Savill 1997).

Suitable species for planting on mine spoils should possess the ability to: (1) grow on poor and dry soils; (2) develop the vegetation cover in a short time and to accumulate biomass rapidly; (3) bind soil to arrest soil erosion and check nutrient loss; and (4) improve the soil organic matter status and soil microbial biomass, thereby enhancing the supply of plant nutrients available. In addition, the species should be economically valuable (Singh et al. 2006). Species with exceptional physiological tolerances to improve site conditions and initiate soil-forming processes means that alien species of *Acacia*, *Alnus*, *Betula*, *Eucalyptus*, *Pinus*, *Salix* and other pioneer trees are frequently employed (Evans 2009a).

Unlike restoration of less severely degraded land, the use of alien tree species remains an acceptable option for mineland revegetation (D'Antonio & Meyerson 2002; Li 2006) if they fulfil a temporary successional role to colonize and ameliorate severely degraded sites and facilitate colonization and eventual dominance by native flora (Seo et al. 2008).

4.2.7 *Alien trees in Short-Rotation Forestry, Short-Rotation Coppice*

Two main drivers have pushed renewable energy production to the top of global agendas: climate change and energy security. Fast-growing poplars and willows can be cultivated in short-rotation forestry (SRF) cycles of 15–18 years, but in short-rotation coppice (SRC) this is reduced further by cut-back/coppicing at 3–5-year intervals (Karp & Shield 2008).

It has been suggested that short-rotation forestry has the potential to deliver greater volumes of biomass from the same land area than alternative biomass crops. Short-rotation forestry is the practice of cultivating fast-growing trees that reach their economically optimum size between eight and 20 years old; each plant produces a single stem that is harvested at around 15 cm diameter. The crops tend to be grown on lower-grade agricultural land, previously forested land or reclaimed land and so do not directly compete with food crops for the most productive agricultural land (McKay 2011).

Of the 330–500 species of willow, the shrub willows (*Salix viminalis* in Europe and *Salix eriocephala* in North America and Canada) are deemed most suitable as bioenergy crops (Kuzovkina et al. 2008). Other species used include *S. dasyclados*, *S. schwerinii*, *S. triandra*, *S. caprea*, *S. daphnoides* and *S. purpurea*, and many varieties are interspecific hybrids. Compared with willows, there are relatively few poplar species that fall into six morphologically and ecologically distinct sections. Of these, Aigeiros (cottonwoods, *Populus nigra*) and *Populus alba* (aspens, white poplars) are of most relevance for bioenergy (Karp & Shield 2008). Many other alien species (including hybrids and genetically modified trees) are used or tested for SRF/SRC, e.g., *Acacia angustissima*, *Gliricidia sepium* and *Leucaena collinsii* in Zambia (Kaonga 2010), *Eucalyptus spp.* and hybrids (e.g., *Eucalyptus grandis* × *E. urophylla* and freeze-tolerant *Eucalyptus* clones), *Platanus occidentalis*, *Pinus taeda*, *Liquidambar styraciflua* in the USA (Hinchee et al. 2009), *Robinia pseudoacacia* in Albania, Italy, Germany, Hungary and Spain (Grünwald et al. 2009; González-García et al. 2011; Rédei et al. 2011a; Kellezi et al. 2012; Ciccarese et al. 2014), *Acacia saligna* in Israel (Eggleton et al. 2007), *Eucalyptus spp.* in UK⁵⁸ (Evans 1980; Leslie et

⁵⁸ The UK has a climate that is not well suited to the majority of eucalypts. However, there is a small number of eucalypt species that can withstand the stresses caused by frozen ground and desiccating winds or sub-zero temperatures that can occur. These species are from more southern latitudes and high-altitude areas of Australia such as, e.g., *Eucalyptus gunnii* (Leslie et al. 2012).

al. 2012) and in China⁵⁹ (Wu et al. 2014), *Acacia* hybrid (*Acacia mangium* × *Acacia auriculiformis*) in Vietnam (Kim et al. 2011), to mention a few.

At the regional scale, significant uncertainties exist and there is a major concern that extensive commercial production with invasive alien trees could have negative effects on biodiversity, in particular in areas of high nature-conservation value. On the contrary, integration of biomass species into agricultural landscapes could stimulate rural economy, thus counteracting to some extent negative impacts of farm abandonment or supporting restoration of degraded land, resulting in improved biodiversity values (Dauber et al. 2010; Bianco et al. 2014).

In Austria⁶⁰ 10 principles for short-rotation forestry systems, from the viewpoint of nature protection and environment, have been declared since 1998 (Trinka 1998). Principle 2 states that “indigenous plants should play an important part, because non-indigenous plants (e.g., *Robinia pseudoacacia* and *Ailanthus altissima*) often show an undesirable tendency to spread”.

4.2.8 Alien trees in agroforestry

Agroforestry systems include both traditional and modern land-use systems where trees are managed together with crops and/or animal production systems in agricultural settings. Agroforestry is practiced in both tropical and temperate regions, for both wood and non-wood products, including food and fibre for improved food and nutritional security (Jama & Zeila 2005). The potential of agroforestry to contribute to sustainable development has been recognized in international policies, including the United Nations Framework Convention on Climate Change (UNFCCC) and the Convention on Biological Diversity (CBD), justifying increased investment in its development (FAO 2013). Agroforestry (or “silvoarable agroforestry”) has traditionally formed important elements of European and Mediterranean landscapes, has the potential to contribute towards sustainable agriculture in Europe in the future, and it is supported by the Common Agricultural Policy (Eichhorn et al. 2006).

Nevertheless, many agroforestry systems, particularly those that depend on tree planting in or near treeless landscapes, rely heavily on alien tree species⁶¹. As is the case in all endeavours based largely on non-native species, problems arise when these alien trees spread from sites of introduction and cultivation to invade areas where their presence is, for various reasons, deemed inappropriate. In some areas, problems caused by the spread of invasive alien agroforestry trees from sites set aside for this land use pose a serious threat to biodiversity that may reduce or negate any biodiversity benefit of the agroforestry enterprise (Richardson et al. 2004).

4.2.9 Alien trees in Mediterranean planted forests and sand dune stabilisation

Planted forests in the Mediterranean have a long history. In mountainous areas, planted forests with introduced coniferous trees were once limited to land at risk from erosion, but these now cover large areas of pastoral land and even agricultural land, either as a result of the naturalisation and establishment of the introduced species (e.g., *Pinus nigra*) or through colonisation of abandoned land. *Pinus radiata* was planted in more than 300,000 ha in old fields, in Spain during the second half of the 20th century, mainly in Atlantic areas. More recently, the species has also been planted in acid soils of the wet Mediterranean area in former agricultural lands restricted to areas with lime-free soils and annual rainfall exceeding 700 mm (Romanyà & Vallejo 2004).

⁵⁹ *E. dunnii*, *E. grandis*, *E. grandis* × *E. camaldulensis*, *E. urophylla* × *E. camaldulensis*, *E. urophylla* × *E. tereticornis*, *E. grandis* × *E. tereticornis*, *E. urophylla* × *E. grandis*.

⁶⁰ The work was done by an interdisciplinary Austrian team of scientists (phytosociologists, landscape ecologists, experts on forestry, on nature protection and on area planning), which was charged with the responsibility by the Austrian Ministry for Science, Transport and Art.

⁶¹ For example, in NW Italy as reported by Sitzia et al. (2013).

In Turkey, afforestation with *P. pinaster* was undertaken by the French for the protection of sand dunes around Terkos Lake in 1880 (Deniz & Yildirim 2014). The first substantial plantings of forest trees in Israel were carried out by the Jewish settlers of Hadera in 1890. They planted Eucalyptus species (mainly *E. camaldulensis*) in an attempt to dry up the nearby swamps and for sand dune stabilisation (Bonneh 2000).

Italian foresters developed successful techniques for stabilising sand dunes, and as a result of their efforts several thousand hectares of dunes were fixed and afforested in Italy and in Libya in the '40s with *Pinus spp.*, *Acacia spp.* and *Eucalyptus spp.* In Libya, beneficial effects were obtained, particularly from the standpoint of protection of highways. Before the Italian Forest Service started its work, the roads from Tripoli to Homs (Lebda) and into the interior were considered unsafe and necessitated many detours (Messines 1952).

BOX 4.2.9.1: Colonisation of Pinus halepensis in Mediterranean habitats: consequences of afforestation, grazing and fire.

Native populations of *Pinus halepensis* in Israel are restricted to the Carmel region and several other mountainous locations. This species was extensively used for afforestation in Israel during the 20th century, and it now constitutes as much as 30% of the planted forests that cover about 100,000 ha within Israel's Mediterranean zone. These forests were planted in a variety of habitat types, some of which are clearly beyond the natural distribution of *P. halepensis* as currently recognized (Lipshitz & Biger 2001). The seed sources that were used for these plantations were mostly alien, and genetically different from the local eastern Mediterranean ecotype (Schiller & Grunwald 1987). There is widespread expansion of *P. halepensis* from plantations into adjacent natural sites, some of which are of high conservation importance (Lavi et al. 2005). This has become an important environmental issue and a topic of ongoing debate among foresters and conservationists. It is clear that the pine expansion is related to the extensive use of *P. halepensis* for afforestation, but the factors that determine the intensity and dynamics of this process are poorly understood (Richardson & Bond 1991; Osem et al. 2011).

4.2.10 Alien trees in arid zones: preventing and combating desertification

Desertification affects millions of the most vulnerable people in Africa, where two thirds of the land cover consists of drylands and deserts. Desertification refers to land degradation in arid, semi-arid and sub-humid areas resulting from factors such as human pressure on fragile eco-systems, deforestation and climate change. Desertification and land degradation have a strong negative impact on the food security and livelihoods of the local communities in Africa's drylands, home to the world's poorest populations (FAO 2014).

Sand encroachment⁶², which has devastating environmental and socio-economic impacts, is another desertification challenge. It reduces arable and grazing land, and diminishes the availability of water resources, threatening the productivity of ecosystems (FAO 2010a).

⁶² Sand encroachment is said to take place when grains of sand are carried by winds and collect on the coast, along water courses and on cultivated or uncultivated land. As the accumulations of sand (dunes) move, they bury villages, roads, oases, crops, market gardens, irrigation channels and dams, thus causing major material and socioeconomic damage. Desertification control programmes must then be implemented in order to counter this very serious situation (FAO 2010b).

The Great Green Wall initiative is a pan-African proposal to “green” the continent from west to east in order to battle desertification. It aims at tackling poverty and the degradation of soils in the Sahel-Saharan region, focusing on a strip of land of 15 km wide and 7,100 km long from Dakar to Djibouti (Dia & Duponnois 2010). The project has faced opposition, despite its stated commitment to combating drought and desertification, and in some case criticised as poorly conceived in terms of both ecological and socio-economic considerations. *Prosopis juliflora*⁶³ is one the species planned to be planted (Dia & Duponnois 2010).

A “Great Green Wall” designed to stop rapidly encroaching deserts and combat climate change is under construction across China. It is a 4,480-km belt of forest across 551 counties and 13 provinces in north-west, central north and north-east China. Part of broader national environment programmes, it is the world’s largest ecological development, and is designed to halt 2,460 km² of land being lost annually to the expanding Gobi Desert due to overgrazing, deforestation and drought. By 2050, the artificial forest is to stretch 400 million ha – covering more than 42 % of China’s landmass. The project began in 1978, and three years later the National People’s Congress, China’s top legislative body, passed a resolution to make it the duty of every citizen above age 11 to plant at least three poplar, eucalypt, larch or other saplings every year (Levin 2005).

4.2.11 Genetically improved and genetically modified alien trees

Diverse biotechnological methods are being intensively pursued to support planted forest with alien trees. These include clonal propagation⁶⁴, interspecific hybridization, the use of a variety of molecular tools to intensify the selection of superior genotypes (DNA fingerprinting, genome mapping, gene identification and genome sequencing) and transformation (Grattapaglia & Kirst 2008; Strauss et al. 2009). However, of this diverse array of technologies, only transformation, defined by the use of direct modification and asexual insertion of DNA into organisms in the laboratory (that is, genetic engineering or modification), engenders attention from the CBD⁶⁵, strong government regulation and controversy over its use, even for research (Strauss et al. 2009).

Research on the strategies and risks of introducing plants with novel traits into natural populations is still in its infancy. Trees are much longer lived and have much longer generation times than annual crops, which makes research more lengthy and difficult. Given the high rates of gene flow found between poplar planted forests and natural populations (Vanden Broeck et al. 2005; Di Fazio et al. 2012), tools for mitigating gene flow should be developed if the introduction of the novel trait is deemed to pose a risk.

Given the diversity of traits, species and environments, a case-by-case approach would seem to be the sensible way to proceed, and this basic approach is officially recognized in the Cartagena Protocol. Annex III/6, under general principles governing risk assessment, states that “risk assessment should be carried out on a case-by-case basis (see also FAO 2010c). The required information may vary in nature and level of

⁶³ Among the 44 recognised *Prosopis* species, *P. glandulosa*, *P. velutina*, *P. juliflora*, and *P. pallida* are considered the most invasive. In Africa, *Prosopis* species are estimated to have invaded over four million ha, threatening crop and range production, desiccating limited water resources, and displacing native flora and fauna (Mwangi & Swallow 2005; Shackleton et al. 2014; Wakie et al. 2014; Shackleton et al. 2015).

⁶⁴ E.g., Rédei et al. (2002, 2011a, b).

⁶⁵ At its 8th Conference of the Parties (COP8) 20-31 March 2006 - Curitiba, Brazil, the Convention on Biological Diversity adopted the Decision VIII/19 “Forest biological diversity: implementation of the programme of work” recommending “Parties to take a precautionary approach when addressing the issue of genetically modified tree”. This Decision recognized “the uncertainties related to the potential environmental and socio-economic impacts, including long-term and transboundary impacts, of genetically modified trees on global forest biological diversity, as well as on the livelihoods of indigenous and local communities, and given the absence of reliable data and of capacity in some countries to undertake risk assessments and to evaluate those potential impacts”. See also COP 9 Decision IX/5 on “Forest biodiversity” 19 - 30 May 2008 - Bonn, Germany, and the EU Directive 2001/18/EC (Directive 2001/18/EC of the European Parliament and of the Council of 12 March 2001 on the deliberate release into the environment of genetically modified organisms and repealing Council Directive 90/220/EEC - Commission Declaration, Official Journal L 106, 17/04/2001 P. 0001 – 0039).

detail from case to case, depending on the living modified organism concerned, its intended use and the likely potential receiving environment.” This principle fits well with the diversity of GM trees (Strauss et al. 2009).

4.3 Invasive alien trees

4.3.1 Generalities and key examples

Many alien trees planted for production or for other purposes have strong direct positive economic impacts on the local and national economies of many countries, but often lead to sharp conflicts of interest when the alien tree species become invasive, and to negative impacts on the ecosystem (Dodet & Collet 2012; van Wilgen & Richardson 2012; Dickie et al. 2014).

Different types of planted forests have provided very important pathways for the introduction and dissemination of invasive alien trees (Wilson et al. 2009; Richardson & Rejmánek 2011; Donaldson et al. 2014). In most cases, alien trees are selected for their adaptability to many habitats, including harsh sites, as well as rapid growth – both features that are shared with invasive alien species (Richardson 1998b).

Evidence has accumulated rapidly around the world on the factors that contribute to invasions of alien trees used in different forms of forestry in the past few decades (Richardson et al. 2014). Importantly, insights on the drivers of such invasions have been shown to be, to some extent and with due care, transferable between regions – regions with recent plantings can learn important lessons from environmentally similar regions in other parts of the world with longer histories of planted forests (Richardson et al. 2015).

Partly because of their large size, but also for other reasons, many alien trees are important ecosystem engineers. Alien tree invasions are among the most costly to manage because invasive alien trees can produce high biomass and they impact on ecosystem services such as water provision where they invade grasslands and shrublands (Richardson 1998a; Le Maitre et al. 2002; van Wilgen & Richardson 2012; Richardson et al. 2014). Many invasive alien tree species cause regime shifts in invaded ecosystems, leading to impacts that ripple across trophic levels (Gaertner et al. 2014).

Alien tree invasions are currently more widespread outside Europe (especially in the southern hemisphere). For this reason, in this Code, many insights are drawn from other regions where appropriate to sketch scenarios and develop principles that are needed to define a code of conduct for Europe and the Mediterranean. In many cases, such insights relate directly to species that are also commonly planted in Europe.

Invasive alien trees can simultaneously bring benefits and cause substantial environmental harm, very often leading to conflicts over how they should be managed. The impacts grow over time as invasions spread, and societal perceptions of the value of invasive alien trees also change as understanding grows and as values shift (Dickie et al. 2014; van Wilgen & Richardson 2014).

The benefits and impacts of invasive alien trees vary in their type and magnitude, depending on the species, their invasive potential, the extent to which they have invaded, and the nature of the invaded environment. The magnitude of benefits and of impacts can be viewed as separate, independent continua, which allows for the classification of species into four broad types (Fig. 1).




Fig. 1 – Types of invasive alien trees based on their relative degree of impact on the environment and the benefits associated with their cultivation and utilisation. Redrawn from van Wilgen & Richardson (2014). The position of any alien tree species within this framework is dynamic.

According to the categorisation proposed by van Wilgen & Richardson (2014), many alien tree species are not invasive, and are either inconsequential, as they have neither substantial impacts nor benefits, or beneficial in cases where they produce useful products, such as wood or fruit, or provide useful ecosystem services, such as sand stabilization or erosion control. It is important to note that the position of any alien tree species within this framework is dynamic. Crucial factors in this regard are the residence time and introduction effort (propagule pressure), but management interventions and changing socio-political conditions can also determine the position of species in this categorization scheme. A few invasive alien tree species provide very little in the way of benefits. Such trees are easily classified as “destructive weeds”, and there is little disagreement about attempts to eradicate or contain such species. Because of the wide variety of uses of trees for humans, there are very few species that can be placed unequivocally in this category. The final type includes species that are both useful and invasive - it is these species that generate controversy and conflict. Finding sustainable solutions to their management is a considerable and escalating problem. Prominent examples in some locations include species in the genera *Acacia*, *Casuarina*, *Eucalyptus*, *Pinus*, *Prosopis* (Wise et al. 2012; Shackleton et al. 2014) and *Salix* (van Wilgen & Richardson 2014).

The number of species falling into this category is increasing rapidly, since the initial benefits of many alien tree species become negated by the impacts when the species become invasive. With an increase in the area planted, the number of species planted and the time since introduction, the number of conflict situations is escalating (van Wilgen & Richardson 2014).

The many benefits of alien trees that are both useful and invasive include: timber production, aesthetic value and appeal, rehabilitation of degraded land, water protection, erosion control and mitigation of sand storms and sand-drift, food for humans, fodder for livestock, carbon sequestration, agroforestry, energy, biodiversity conservation (van Wilgen & Richardson 2014), facilitation of indigenous tree species regeneration, recreation and landscape amenity. Responsible management of planted forests has a critical benefit to natural forest ecosystems as it reduces the pressures on indigenous forest for forest products and allows them to be designated for other protective and conservation purposes (FAO 2010c; Mead 2013).

The role of planted forests of alien trees in biodiversity conservation is now better understood (Brockerhoff et al. 2008). For example, in the fragmented forest landscape of Europe, planted forests comprise an increasingly important fraction of the matrix surrounding natural forest fragments. Planted forests may provide habitat for forest biodiversity, which may enhance landscape connectivity and regional biodiversity (Bremer & Farley 2010).

Quine and Humphrey (2009) compared the species richness of a range of different taxonomic groups (lichens, bryophytes, fungi, vascular plants, invertebrates and songbirds) in alien and native forest stands of differing structural stages in northern and southern Britain. In terms of overall native species-richness no significant difference between the alien and the native stands was detected, but lichen species richness was much lower in the alien stands compared to the native stands, whereas bryophyte and fungal species richness was proportionately higher in the alien stands. They concluded that emergent ecosystems of alien conifer species are not irrelevant to biodiversity. Where already well-established they can provide habitat for native species particularly if native woodland is scarce and biodiversity restoration is an immediate priority (Quine & Humphrey 2009).

On the contrary, Calviño-Cancela (2013) described a paucity of birds in planted forests of *Eucalyptus globulus* in Galicia, Spain, compared with the diverse and abundant avifauna in planted forests of pines. The alien status and taxonomic isolation of *E. globulus* in the region, together with specific features of its leaves and bark, may explain the low suitability of eucalypt planted forests, by limiting the presence of phytophagous insects and thus the availability of prey for birds (Calviño-Cancela 2013).

Invasive alien trees escaping from planted forests may affect invaded ecosystems and ecosystem functioning and services in many ways. Many act as **transformers** (ecosystem engineers), i.e. they are invasive alien trees that reach very high densities and substantially increase biomass or change the type and arrangement of above-ground material. In some cases, the impacts of tree invasions are obvious, fast and dramatic. Some invasions have radically transformed entire ecosystems.

For example, invasive alien *Acacia* and *Pinus* species have rapidly transformed species-rich fynbos shrublands in South Africa and sand-dune vegetation in Portugal into species-poor, forests or woodlands dominated by alien species and with markedly changed biodiversity and ecosystem functions. Invasion of *Melaleuca quinquenervia* in Florida's Everglades⁶⁶ has changed large areas of open grassy marshes to closed-canopy swamp-forests.

The net effect of invasive alien trees is determined by the product of the per-capita effect, the abundance they achieve (reflected by numbers of stems per area, or the total biomass added), and their geographical range (Parker et al. 1999). Some invasive alien trees have no obvious impacts, some have localized impacts, while others cause massive ecosystem-level transformations. Some of the most prominent effects of invasive alien trees in different parts of the world are summarized below, following the impact categories of Richardson et al. (2000). Full details and references for the examples given are provided in Richardson & Rejmánek (2011).

Excessive users of resources: Many invasive alien trees invade riparian ecosystems where they achieve dominance and huge abundance and thus consume more water than would the native species that normally frequent these ecosystems. The impact is due primarily to increased biomass and therefore increased water use. Prominent examples are *Tamarix* spp.⁶⁷ in SW North America (Stromberg et al. 2007) and *Acacia* species, notably *A. mearnsii*, in South Africa (Dye & Caren 2004) and in South Sardinia (Italy).

Donors of limiting resources: Many invasive alien species of woody legumes impact invaded ecosystems primarily via their addition of nitrogen (natural fertilization). Well-studied examples are *Morella faya* which doubles canopy nitrogen as it replaces native forest species in Hawaii, the Australian

⁶⁶ *Melaleuca quinquenervia*, introduced to Florida in the late 1800s as an ornamental and for other purposes (drying up wetlands), is now the most prominent of 60 non-native plant species invading many natural wetland and upland areas in Florida, including the Everglades, a United Nations World Heritage site and UNESCO Man and Biosphere Reserve. *Melaleuca* invasion has determined displacement of native species, reduction in wildlife habitat value, alteration in hydrology, modification of soil resources, changes in fire regimes (Mazzotti et al. 2014).

⁶⁷ Tamarisk taxa (*Tamarix ramosissima*, *T. chinensis* and their hybrids) were introduced to the United States from Asia in the late 1800s for the control of soil erosion and landscaping purposes. They are now the third most prevalent alien tree riparian taxon in the western United States (Friedman et al. 2005).

Acacia species in South Africa and in the Mediterranean, and *R. pseudoacacia* which increases the soil nitrogen pools in nitrogen-poor soils in Europe (e.g., Sitzia et al. 2012; Cierjacks et al. 2013; González-Muñoz et al. 2013). *A. altissima* increases the availability of mineral nitrogen under its canopy due to the large amounts of fast-decomposing litter that it produces (González-Muñoz et al. 2013; Medina-Villar et al. 2015).

Fire promoters/suppressors: The best-studied example of an invasive alien tree that brings fire to a previously fire-free system is that of *Melaleuca quinquenervia* invasions of wetland habitats in Florida, USA, where a massive increase in flammable material leads to very intense fires. Examples of where alien tree and shrub invasions have suppressed fire frequency are *Mimosa pigra* in northern Australia and *Triadica sebifera* and *Schinus terebinthifolius* in North America; in all cases alien tree invasions result in reduced horizontal continuity of fuel which reduces fire frequency and intensity (Brooks et al. 2004).

Sand stabilizers: Australian *Acacia* species have been widely planted along coastal dunes in several parts of the world to stabilize sand movement. Planted and self-sown stands of species like *A. cyclops* perform this function very well; in some areas of South Africa dune stabilization has resulted in massive beach erosion. In the Portuguese dune ecosystems, *Acacia longifolia* and *A. saligna* are among the most aggressive invasive alien tree species. These alien woody legumes were planted at the beginning of the last century to curb sand erosion but have now proliferated, often associated to fire events, causing significant ecological impacts. Long-term occupation by *A. longifolia* significantly altered the soil properties with increased levels of organic C, total N and exchangeable cations resulting in higher microbial biomass, basal respiration, and b-glucosaminidase activity (Marchante 2001; Marchante et al. 2003; Marchante et al. 2008). The replacement of drought tolerant native species by the water spending invader, *A. longifolia*, may have serious implications for ecosystem functioning, especially during the prolonged drought periods predicted to occur in Portugal in the future (Rascher et al. 2011).

Colonisers of intertidal mudflats/sediment stabilizers: Red mangrove (*Rhizophora mangle*) was introduced to Hawai'i in 1902 to control runoff from upstream agriculture. Other species of alien mangrove have been introduced to Hawai'i, but *R. mangle* is the most successful, occupying coastal habitats throughout the main Hawaiian Islands, including estuarine fishpond sites developed for aquaculture by native Hawaiians as early as 1000 C.E. (Siple & Donahue 2013).

In their native range, mangroves are ecosystem engineers, strongly modifying their environment and providing important ecosystem services, including shoreline protection, entrapment of heavy metals, sediment stabilization, litterfall subsidy, and nursery grounds (Siple & Donahue 2013). In their introduced range, these potential ecosystem services must be weighed against impacts on native ecosystems: In Hawai'i, alien mangroves create habitats dramatically distinct from the sandflats inhabited by the few native coastal macrophytes, transforming nearshore sandy habitat into heavily vegetated areas protected from oceanic erosion with low water velocity, high sedimentation rates, and anoxic sediments. Alien mangrove forests provide habitat for alien species, including burrowing predators, which can exert top-down effects on benthic communities (Siple & Donahue 2013).

Litter accumulators: The North American *Pinus strobus* invades both natural *P. sylvestris* forests and planted forests of the latter species in sandstone areas of the Czech Republic. *Pinus strobus* produces greater quantities of more slowly decomposing litter than its native congener which has a major effect on soil acidity. Under such conditions, *P. strobus* regenerates better than *P. sylvestris* which contributes to its success as an alien invader (Pyšek & Prach 2003). In Central Europe, many sandstone areas are protected for their unique environment, and large-scale regeneration of this invasive alien tree species is of concern (Hadincová et al. 2007; Mandák et al. 2013).

Invasive alien tree species can **hybridise and introgress** if the species have taxonomical close relatives in the native flora. This can be desirable for a better adaptation to changing climatic conditions and to meet human needs in the renewable material “wood”, while undesirable from a conservation point of view (Rhymer & Simberloff 1996; Smulders et al. 2008; Felton et al. 2013; Kjær et al. 2014), especially if the native species are rare⁶⁸ in number compared to planted individuals of the invasive alien tree.

Many invasive alien trees qualify as “**transformers**” sensu Richardson et al. (2000). Well-studied examples of are Australian *Acacia* species (in Chile, Portugal, South Africa), *Cinchona pubescens* (Galapagos islands), *Ligustrum robustum* var. *walkeri* (La Réunion Island), *Melaleuca quinquenervia* (Florida, USA), *Miconia calvescens* (Tahiti), *Mimosa pigra* (northern Australia & Zambia), *Morella faya* (Hawaii), *Pinus pinaster* (South Africa), and *Triadica sebifera* (North America). On the contrary, in France, *Pinus pinaster* transformed a poor scrubby acid sandy soil into a productive forest of 1 million hectares that fulfills many economic, social and environmental functions (pers. comm. MAAF 2016).

Besides the effects mentioned above that are attributable to effects on physical resources either due to large size and biomass or impacts on resource availability, many alien tree and shrub invasions affect resident biota in more subtle ways. An important category of impacts for invasive alien tree species is the alteration of habitat for other organisms. A few examples from different parts of the world illustrate the very wide range of changes that invasive alien trees can cause. In Hawaii, the spread of alien mangroves has led to habitat loss for wetland birds (Allen 1998; Siple & Donahue 2013). The new mangrove habitats also provide refugia for shorebird predators, including invasive rats (*Rattus* spp.) and mongooses (*Herpestes* spp.), and alien marine species such as the mangrove crab (*Scylla serrata*). Emergent roots of invasive *Rhizophora mangle* are also colonized by various introduced barnacles and sponges, thus altering the structure of macrofaunal communities. Many invasive alien trees have a major impact by creating impenetrable thorny thickets that limit the passage of animals (e.g., *Caesalpinia decapetala*, *Mimosa pigra*, and *Prosopis* spp.). *Annona glabra* invades Australian estuaries and chokes mangrove swamps, where its seedlings carpet the banks and prevent other species from germinating or surviving. Invasion of these riparian zones by willows (*Salix* species) decreased food resources and altered habitat, reducing native bird diversity and disrupting connectivity of the riparian zone. On the island of Sao Miguel in the Azores archipelago, invasion of the native forest by the alien tree *Pittosporum undulatum* and other alien species led to a marked reduction in structural complexity and an impoverished flora. This led to a reduction in insect biomass, due to the replacement of large insects on native plants with small insects on alien plants. This appears to have far-reaching negative consequences for ecosystem stability (Heleno et al. 2009).

Besides the diverse ecological effects discussed above (many of which are associated with modification of ecosystems), alien tree invasions have many complex effects on human livelihoods, both positive and negative. These have been clearly documented in South Africa (especially for Australian acacias) and Papua New Guinea (due to invasion of *Piper aduncum*). *Prosopis* invasions in sub-Saharan Africa have led to considerable rangeland degradation, causing many problems for human societies, especially those relying on subsistence agriculture (e.g., Mwangi & Swallow 2005; Shackleton et al. 2014). Alien tree invasions have huge financial costs in many regions.

In Britain, several alien trees have become culturally naturalised (Mabey 1996; Peterken 2001) causing a change in the perception of nature. *F. sylvatica* in northern and western Britain is widely accepted by the general public as a native. *A. pseudoplatanus* is regarded as traditional by remote farm buildings in Wales and northern England. *P. sylvestris* is seen as a natural part of the scenery in southern heathlands (Peterken 2001).

⁶⁸ E.g. *Abies nebrodensis*, or Sicilian fir, is an endemic species of Sicily, Italy, growing on the Madonie range at 1700–1900 m above sea level. It is a highly-endangered species (Council of Europe 1977), comprising a single relict population of approximately 30 adult trees spread over an area of 150 ha (Ducci 2014).

Alien tree invasions into grasslands and shrublands convert many unique vegetation formations into virtual monocultures of alien trees. Macdonald et al. (1988) recognized the analogy between invasion of *Cinchona pubescens* into shrubby highland communities of the Santa Cruz Island, Galapagos, and invasion of alien pines and acacias into fynbos shrublands in South Africa. These alien tree invasions are key contributors to the degradation of such ecosystems over much of their extent.

Picea sitchensis originates from the west coast of North America and is imported to Norway mainly as a production species for planted forests, although it is also used for shelter belts. Historically, the species has also been imported for research purposes, but such importations have now ceased. *Picea sitchensis* has been established in Norway since the '50s and is the most important alien tree species in terms of extent of planting. The species is mainly planted in coastal areas in heaths, grazed blueberry forest and small fern forest. It is considered an invasive alien primarily in these types of habitats where it is expected to spread further (Gederaas et al. 2012).

Box 4.3.1.1 Invasive alien Acacia species.

Invasive alien *Acacia* species, like many other invasive alien species, have many types of impacts including some that interact synergistically. Alien *Acacia* species can induce simultaneous changes in the above- and below-ground communities, microclimates, soil moisture regimes and soil nutrient levels (Marchante et al. 2003, 2008b; Yelenik et al. 2004; Werner et al. 2010; Gaertner et al. 2011). Many changes are directly attributable to key traits of *Acacia* species: their rapid growth rates and ability to out-compete native plants (Morris et al. 2011); their capacity to accumulate high biomass; large, persistent seed banks; and their capacity to fix nitrogen (Yelenik et al. 2007). These features enable them to dominate competitive interactions with native species. Many of the abiotic changes and biotic responses to them are tightly linked and may advance simultaneously rather than sequentially (Hobbs et al. 2009), as does the progression from structural to functional impacts (Le Maitre et al. 2011). The impacts of invasive alien Australian acacias on biodiversity and ecosystem properties and functions also affect the delivery of ecosystem services and the benefits that society derives from them. Affected ecosystem services include: supporting services (e.g. soil formation); regulatory services (e.g. water flow and nutrient cycling); production services (e.g. food and fibre); and cultural or life-enhancing services (e.g. recreation or educational opportunities to sustain human well-being) (Le Maitre et al. 2011).

Finally, past experience of introducing tree species from other countries shows that one of the possible negative outcomes could be the failure of the introduced tree to grow successfully (Engelmark et al. 2001). Alien tree species are widely used in planted forests for their high productivity and performance compared to native trees. However, these advantages may be compromised, as for any tree species, by herbivore damage, insects and microbial pathogens, which are introduced accidentally and/or have adapted to new host trees (Branco et al. 2014; Wingfield et al. 2015).

4.4 International initiatives and legislation on invasive alien species and invasive alien trees

Many international instruments refer to invasive alien species that may have undesired environmental or economic impacts. These range from legally binding treaties to non-binding technical guidance focused on particular species or pathways. The main international regulations concerning invasive alien species are given in the following with specific reference to invasive alien trees and planted forests. The information here provided is intended to provide support to the principle “be aware of regulations” of the present Code.

4.4.1 The Convention on Biological Diversity

The Convention on Biological Diversity (CBD), negotiated under the auspices of the United Nations Environment Programme (UNEP), was adopted in 1992 and entered into force in 1993. Its aims are the conservation of biological diversity, the sustainable use of biological resources, and the fair and equitable sharing of benefits arising from the use of genetic resources (Secretariat of the Convention on Biological

Diversity 2001a, 2001b). CBD requires Parties “as far as possible and as appropriate (to) prevent the introduction of, control or eradicate those alien species which threaten ecosystems, habitats or species” under the provision of Article 8(h).

Given this mandate, the Convention’s member governments who together constitute the Conference of the Parties (COP) to the Convention made numerous decisions with respect to alien species, many of which are directly relevant to the management of (invasive) alien tree species. In particular, the COP 11 Decision XI/19⁶⁹ states that “when designing, implementing and monitoring afforestation, reforestation and forest restoration activities for climate change mitigation, consider conservation of biodiversity and ecosystem services through, for example: (i) Converting only land of low biodiversity value or ecosystems largely composed of non-native species, and preferably degraded ecosystems; (ii) Prioritizing, whenever feasible, local and acclimated native tree species when selecting species for planting; (iii) Avoiding invasive alien species; (iv) Preventing net reduction of carbon stocks in all organic carbon pools; (v) Strategically locating afforestation activities within the landscape to enhance connectivity and increase the provision of ecosystem services within forest areas”.

4.4.2 *The Council of Europe and the Bern Convention*

The Council of Europe⁷⁰ promotes actions (without any regulatory mandate), to avoid the intentional introduction and spread of alien species, to prevent accidental introductions and to build an information system on invasive alien species (IAS). In 1984, the Committee of Ministers of the Council of Europe adopted a recommendation to that effect. Also, the Bern Convention (Convention on the Conservation of European Wildlife and Natural Habitats), the main Council of Europe treaty in the field of biodiversity conservation, requires its Contracting Parties “to strictly control the introduction of non-native species⁷¹”.

In 2003, the Bern Convention adopted the European Strategy on Invasive Alien Species (Genovesi & Shine 2004), aimed at providing precise guidance to European governments on IAS issues. The Strategy identifies European priorities and key actions, promotes awareness and information on IAS, strengthening of national and regional capacities to deal with IAS issues, taking of prevention measures and supports remedial responses such as reducing adverse impacts of IAS, recovering species and natural habitats affected. National strategies have been drafted and implemented by many of the Parties following the priorities set in the European Strategy. Noteworthy, many Recommendations on IAS have been adopted by the Standing Committee since 1997⁷².

⁶⁹ COP 11 Decision XI/19, Hyderabad, India, 8-19 October 2002 - “Biodiversity and climate change related issues: advice on the application of relevant safeguards for biodiversity with regard to policy approaches and positive incentives on issues relating to reducing emissions from deforestation and forest degradation in developing countries; and the role of conservation, sustainable management of forests and enhancement of forest carbon stocks in developing countries”. Cf. also Secretariat of the Convention on Biological Diversity (2002) (<https://www.cbd.int/doc/publications/cbd-ts-07.pdf>). The section on “unsustainable forest management” reports case studies on *Leucaena leucocephala*, *Miconia calvescens*, *Spathodea campanulata* and *Cordia alliodora* impacts.

⁷⁰ The Council of Europe includes 47 member states, 28 of which are members of the European Union. (<http://www.coe.int/en/web/portal/home>).

⁷¹ In Article 11, paragraph 2.b of the Convention, each Contracting Party undertakes to strictly control the introduction of non-native species.

⁷² Recommendation No. 57 (1997) on the Introduction of Organisms belonging to Non-Native Species into the Environment; Recommendation No. 91 (2002) on Invasive Alien Species that threaten biological diversity in Islands and geographically and evolutionary isolated ecosystems; Recommendation No. 77 (1999) on the eradication of non-native terrestrial vertebrates; Recommendation No. 99 (2003) on the European Strategy on Invasive Alien Species, which recommends that Contracting Parties: draw up and implement national strategies on invasive alien species taking into account the European Strategy on Invasive Alien Species. And co-operate, as appropriate, with other Contracting Parties and Observer States in prevention, mitigation and eradication or containment of alien species; Recommendation No. 134 (2008) of the Standing Committee, adopted on 27 November 2008, on the European Code of Conduct on Horticulture and Invasive Alien Plants; Recommendation No 141 (2009) of the Standing Committee, adopted on 26 November 2009, on potentially invasive alien plants being used as biofuel crops.

4.4.3 *The International Plant Protection Convention*

The International Plant Protection Convention (IPPC), which has existed since the 1950s, aims to prevent the introduction and spread of plant pests. National plant protection services and the governing body of the IPPC, the Interim Commission on Phytosanitary Measures (ICPM), recognized that the aim of the CBD to prevent the introduction of alien species corresponds in large measure to the aim of the IPPC. Since 1999, the ICPM has been actively engaged in clarifying its role in regard to invasive alien species that are plant pests. In 2001, it determined that such species should be considered quarantine pests and should be subjected to measures according to IPPC provisions. The ICPM also decided that IPPC standards should be reviewed to ensure that they adequately address environmental risks of plant pests. In 2003, the ICPM adopted supplements to two of the international standards for phytosanitary measures (namely Glossary of phytosanitary terms and Pest risk analysis for quarantine pests). These supplements elaborated on environmental considerations. To avoid conflicting developments within the IPPC and the CBD regarding invasive alien species and plant pests (Lopian 2005; Brunel et al. 2009).

4.4.4 *The European and Mediterranean Plant Protection Organisation (EPPO)*

The European and Mediterranean Plant Protection Organization (EPPO) is an intergovernmental organization responsible for European cooperation in plant health. Nearly all countries of the European and Mediterranean region are members. EPPO's objectives are to protect plants, to develop international strategies against the introduction and spread of dangerous pests and to promote safe and effective control methods. It is developing a cooperative Europe-wide strategy to protect the EPPO region against invasive alien plants and created in 2002 a Panel on Invasive Alien Species which was charged with identifying invasive plant species that may present a risk to the EPPO region and proposing measures to prevent their introduction and spread and recommendations on ways to eradicate, suppress and contain invasive alien species that have already been introduced (Brunel et al. 2009).

The Panel has established the EPPO List of Invasive Alien Plants which can be considered as a list of priorities. The alien trees *Acacia dealbata*, *Ailanthus altissima* and *Prunus serotina* are listed in the EPPO list of invasive alien plants⁷³.

EPPO publishes standards and guidelines, the EPPO Bulletin and the EPPO reporting systems, as valuable sources of information on invasive alien species. During 2016, a three-day workshop was held at the headquarters EPPO, with the purpose of prioritising a list of invasive alien plants for risk assessment under the LIFE⁷⁴ funded project 'Mitigating the threat of invasive alien plants in the EU through pest risk

Recommendation No. 142 (2009) the Standing Committee, adopted on 26 November 2009, interpreting the CBD definition of invasive alien species to take into account climate change. It recommends Contracting Parties to the Convention and invites Observer States to: "interpret the term "alien species" for the purpose of the implementation of the European Strategy on Invasive Alien Species as not including native species naturally extending their range in response to climate change"; Recommendation No. 160 (2012) of the Standing Committee, adopted on 30 November 2012, on the European Code of Conduct for Botanic Gardens on Invasive Alien Species; Recommendation No. 179 (2015) of the Standing Committee - Strasbourg, 1 December - 4 December 2015, on action to promote and complement the implementation of EU Regulation 1143/2014 on invasive alien species. (<http://www.coe.int/en/web/bern-convention/recommendations-on-ias>).

⁷³ The plants listed have been identified by the EPPO Panel as being absent or present in the EPPO region; as having a high potential for spread; as posing an important threat to plant health and/or the environment and biodiversity; and eventually as having other detrimental social impacts in the EPPO region. Because a large number of invasive alien plants are already present in the EPPO region, priorities were set in order to select those species considered to pose the greatest threat to species and ecosystems in the EPPO region. EPPO therefore strongly recommends countries endangered by these species to take measures to prevent their introduction and spread, or to manage unwanted populations (for example with publicity, restrictions on sale and planting, and control measures). This List is constantly being reviewed by the Panel (new species can be added and others removed). The list is not meant to be exhaustive but to focus on the main risks (http://www.eppo.int/INVASIVE_PLANTS/ias_lists.htm).

⁷⁴ LIFE is the EU's financial instrument supporting environmental, nature conservation and climate action projects throughout the EU (<http://ec.europa.eu/environment/life/>).

analysis to support the EU Regulation 1143/2014⁷ (LIFE15 PRE FR 001) (see, www.IAP-risk.eu). The workshop was comprised of experts from the EPPO Panel on Invasive Alien Plants, the NERC Centre for Ecology and Hydrology and the EPPO Secretariat. A group of alien tree species were identified as having a high priority for a risk assessment, i.e. *Acacia dealbata*, *Cinnamomum camphora*, *Hakea sericea*, *Prosopis juliflora* and *Sapium sebiferum*.

4.4.5 *The Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES)*

CITES, which primarily addresses trade in endangered species, can prevent or better regulate the transfer of endangered species that may be invasive alien. It has three different levels of protection for species, reported as Appendices⁷⁵. Although there are literally thousands of plant species protected under CITES, only a portion of these species are trees, and of the included tree species, only a relatively small portion of them are actually used as lumber. *Araucaria araucana*⁷⁶ and *Dalbergia nigra*⁷⁷ are included in Appendix I.

4.4.6 *Sustainable forest management and forest certification*

The idea of having international common guidelines in the forest sector dates back to 1992. The Forest Principles is the informal name given to the “Non-Legally Binding Authoritative Statement of Principles for a Global Consensus on the Management, Conservation and Sustainable Development of All Types of Forests”, a document produced at the United Nations Conference on Environment and Development (UNCED), informally known as the Earth Summit. It is a non-legally binding document that makes several recommendations for conservation and sustainable development of forestry. Since a statement is, by its nature, non-legally binding, the title's inclusion of these words shows that this non-binding aspect needed extra emphasis, demonstrating the great divergence of views during the UNCED negotiations in 1992 (Ruis 2001). These principles should apply to all types of forests, both natural and planted, in all geographical regions and climatic zones and, importantly, principle 6 (a) states that “all

⁷⁵ Appendix I: This appendix represents species that are in the most danger and are considered to be threatened with extinction, and are consequently the most restricted in international trade. Appendix II: This appendix contains species that are at risk in the wild, but not necessarily threatened with extinction. Species in this appendix are closely regulated, but are typically not as restricted as Appendix I. Appendix III: This appendix contains species that a certain country (called a “party” within CITES), has voluntarily requested to be regulated in order to help preserve the species in question. Appendix III species regulation is only applicable for the specific party that has requested its inclusion, and is therefore much less restrictive than Appendix I or II. CITES is implemented in the EU through the Wildlife Trade Regulations. Currently these are Council Regulation 338/97/EC on the protection of species of wild fauna and flora by regulating trade therein (the Basic Regulation) and Commission Regulation 865/2006/EC laying down detailed rules concerning the implementation of Council Regulation 338/97/EC (the Implementing Regulation). Suspension regulations including 997/2010/EC (5 November 2010) and Regulation 359/2009/EC (30 April 2009) suspend the introduction into the Community of certain species from certain countries.

⁷⁶ *Araucaria araucana* has been widely planted as a specimen tree in temperate areas all over the world, but there are virtually no planted forests. A small-scale plantation was established in southwest Scotland in 1916 (Williams & Winn 1977; Premoli et al. 2013). Endangered for IUCN, *Araucaria araucana* is listed on Appendix I of CITES which strictly regulates the trade in its timber and seeds (CITES 2014: <http://www.iucnredlist.org/details/31355/0>).

⁷⁷ *Dalbergia nigra* (Vell.) Allemao ex Benth, known as the Brazilian rosewood or jacarandá-da-Bahia, is a tree species endemic to the central Atlantic Forest in Brazil. This species produces a high-quality wood that is highly valued for the manufacture of musical instruments and fine furniture, thus resulting in its overcutting since the colonization of Brazil. *D. nigra* is a threatened tree that is in the “Endangered” category due to its over-exploration, the absence of replacement plantations and the deforestation of the Atlantic Forest - (IUCN). *D. nigra* is extremely rare in nature, and its international trade has been prohibited since the 1990s by the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES 2008, Appendix I, II and III to the Convention on International Trade in Endangered Species of Wild Fauna and Flora. US Fish and Wildlife Service: Washington; Ribeiro et al. 2011; Taylor et al. 2012). Further information can be found in the Resolution Conf. 13.10 (Thirteenth meeting of the Conference of the Parties - <http://www.cites.org/eng/res/13/13-10.php>).

types of forests play an important role in meeting energy requirements through the provision of a renewable source of bio-energy, and that the potential contribution of plantations of both indigenous and introduced species for the provision of both fuel and industrial wood should be recognized”.

The FOREST EUROPE process (Ministerial Conferences on the Protection of Forests in Europe, MCPFE), a pan-European voluntary political process for dialogue and cooperation on forest policies in Europe, was started by the Strasbourg Conference in 1990 and the Forest Principles were adopted and incorporated into the agenda by Helsinki Conference in 1993. The concept of sustainable forest management (SFM) was defined as: "The stewardship and use of forest lands in a way and at a rate that maintains their productivity, biodiversity, productivity, regeneration capacity, vitality and their potential to fulfil now and in the future relevant ecological, economic and social functions at local, national and global levels and that does not cause damage to other ecosystems" (European Commission 2016; FOREST EUROPE 2016).

FOREST EUROPE develops common strategies for its 47 signatories (46 European countries and the European Union) on how to protect and sustainably manage their forests. Since 1990, the collaboration of the ministers responsible for forests in Europe has had a great economic, environmental and social impact on the national and international level. FOREST EUROPE has led to achievements such as the guidelines, criteria and indicators for sustainable forest management (FOREST EUROPE 2016).

The first two set of guidelines, “General guidelines for sustainable forest management” and “General guidelines for conservation of biological diversity of forests in Europe” were developed in Helsinki in 1993 (Resolution H1 and H2 respectively). They were elaborated as general political guiding principles to be implemented in an integrated manner to be reflected in national guidelines and local technical solutions.

In the first part of the Resolution H1 “general guidelines”, principle 9 states that “*Native species and local provenances* should be preferred where appropriate. The use of species, provenances, varieties or ecotypes outside their natural range should be discouraged where their introduction would endanger important/valuable indigenous ecosystems, flora and fauna. *Introduced species* may be used when their potential negative impacts have been assessed and evaluated over sufficient time, and where they provide more benefits than do indigenous ones in terms of wood production and other functions. Whenever introduced species are used to replace local ecosystems, sufficient action should be taken at the same time to conserve native flora and fauna”.

The “Pan-European Operational Guidelines for Sustainable Forest Management” endorsed at Lisbon Ministerial Conference in 1998, were further elaborated to translating international commitments to the level of forest management practices and planning. They are directly based on Resolutions H1 and H2, and follow the structure of the six pan-European criteria that were identified as the core elements of sustainable forest management. They are divided into/addressing “Forest Management Planning” and “Forest Management Practices”, focusing on basic ecological, economical and social requirements for sustainable forest management within each criterion. The Criterion no. 4 is titled “Maintenance, conservation and appropriate enhancement of biological diversity in forest ecosystems” and at 4.2 (b) states that “for reforestation and afforestation, origins of *native species and local provenances* that are well adapted to site conditions should be preferred, where appropriate. Only those *introduced species, provenances or varieties* should be used whose impacts on the ecosystem and on the genetic integrity of native species and local provenances have been evaluated, and if negative impacts can be avoided or minimised”.

FOREST EUROPE has also developed, in cooperation with the Environment for Europe/Pan European Biological and Landscape Diversity Strategy, the “**Pan-European Guidelines for Afforestation and Reforestation**” with a special focus on the provisions of the UNFCCC. The Guidelines, agreed in 2008, recognize the role of sustainable forest management in climate change mitigations. They form a set of recommendations for voluntary use by national authorities and other bodies and stakeholders relevant to implement economically viable, environmentally sound and socially equitable afforestation and

reforestation programmes and projects. In the section titled “Ecological Guidelines”, guideline 19 affirms that “*native tree species*, provenance and varieties or ecotypes that are well adapted to site conditions should be used for afforestation and reforestation where appropriate”; guideline 20 that “the need to consider adaptation to climate change should be taken into account when choosing species, provenances and varieties for afforestation and reforestation” and guideline 21 that “species, provenances, varieties or ecotypes *outside their natural range* should only be used where their introduction would not endanger important and/or valuable indigenous ecosystems, flora and fauna. Those that are likely to be invasive should be avoided using the CBD Guiding Principle for the Prevention, Introduction, and Mitigation of Impacts of Alien Species That Threaten Ecosystems, Habitats or Species”. In addition, guideline 22 states that “a precautionary approach should be taken to the use of *genetically modified trees*. Ecological socio-economic and cultural impacts, including long term effects should be analysed and a thorough, comprehensive and transparent risk assessment should be completed in accordance with the Cartagena Protocol on Biosafety. In this context, the potential impacts of genetically modified trees on native gene pools should be fully considered”.

Therefore, standards, guidelines, criteria and indicators for sustainable forest management have been developed over the past few decades by intergovernmental processes, international organizations⁷⁸, certification schemes (e.g. FSC, PEFC) (Masiero et al. 2015) and national governments. These apply to all forests, including planted forests, and have resulted in forestry being recognised as an essentially sustainable land-use and essential to combatting climate change by storing carbon and preventing deforestation. Activity was increased considerably after the Statement of Principles for the Sustainable Management of Forests was adopted in 1992 at the Earth Summit in Rio in response to global concerns about deforestation and the unsustainable exploitation of natural forests (Stupak et al. 2011). At European level, the 46 signatories of the Ministerial Conference on the Protection of Forests in Europe agreed a definition of sustainable forest management in a Ministerial Process dating from 1990 and have developed and refined a set of criteria and indicators. These are regularly updated and adapted to new challenges⁷⁹. In addition, the International Tropical Timber Organization ITTO has developed and is revising criteria and indicators for sustainable forest management since the early 1990’s⁸⁰. Planted forests can take over many, though not all, functions that indigenous forest provide (FAO 2010c) and contribute enormously to economic benefits whilst relieving the pressure on indigenous forests.

Forest certification is a voluntary sustainable forest management tool that aims to promote the sustainable management and conservation of forest ecosystems by adding market value to products generated according to environmental and socio-economic principles (Cashore et al. 2004; Auld et al. 2008; Gomez-Zamalloa et al. 2011; Meidinger 2011; Dias et al. 2013). It is based on third-party auditing of compliance with environmental and socio-economic standards, developed by governmental actors, environmental non-governmental organizations, industry associations, and social groups through participatory public processes. Forest certification relies on the willingness of a growing number of consumers to pay more for sustainably generated products and it aims to reward forest managers that follow sustainable forest management practices (Auld et al. 2008; Brown et al. 2001; Suzuki & Olson 2008).

Two forest certification systems dominate globally: The Forest Stewardship Council (FSC⁸¹) and the Programme for the Endorsement of Forest Certification (PEFC⁸²). FSC is a global forest certification

⁷⁸ Cf., New Generation Plantations (NGP), 2014. New generation plantations: review 2014 (<http://newgenerationplantations.org/multimedia/file/12b486cb-ea24-11e3-9f9e-005056986313>).

⁷⁹ Forest Europe http://www.foresteurope.org/sfm_criteria/criteria

⁸⁰ ITTO http://www.itto.int/sustainable_forest_management/

⁸¹ Forest Stewardship Council (FSC), a not-for-profit international organization established in the early 1990s to promote the responsible management of the world’s forests. FSC has a framework of globally applicable Principle and Criteria (FSC Std 01-001). For each of the Criteria indicators are developed for “local” certification – specific to the national legal, ... circumstances. These are the national standards (i.e. the text of the P&C is in all standards exactly the same – only the indicators might differ from country to country). In some countries these national

scheme. PEFC is an umbrella organization that endorses national schemes, some of which were developed within the PEFC framework, while others existed as independent schemes for several years before PEFC was formed, e.g., American Tree Farm System (ATFS), Sustainable Forestry Initiative (SFI) and Canadian Standards Association (CSA) (Stupak et al. 2011).

FSC certification was created in 1993 to “promote environmentally appropriate, socially beneficial, and economically viable management of the world’s forests” (Auld et al. 2008; <https://ic.fsc.org/index.htm>). FSC certification⁸³ comprises 10 principles and 70 criteria that cover environmental, social and economic aspects of forest management. The standard uses the CBD definition of alien species and criterion 10.3 (Principle 10 “Implementation of Management Activities”) states that “The Organization⁸⁴ shall only use alien species⁸⁵ when knowledge and/or experience have shown that any invasive impacts can be controlled and effective mitigation measures are in place”.

Canada has the largest area of third-party independently certified forests (CSA, FSC, SFI) in the world. As of 2011, more than 151 million ha of Canadian forests were certified, which represents 42% of the world’s forests under certification. The UK was the first country in the world to have all its state forests independently certified. Sweden was the first country to introduce a national system of certification based on FSC standards. The country has, as a result, a disproportionate part of the FSC portfolio even though its share of the total certified area has declined in recent years; from about 30 % of the world's FSC-certified forests in the beginning of the century (Boström 2003) to 12 % (Regional Totals: Forest Management Certifications, 8 October 2007, <http://www.fsc-sverige.org>) (Schlyter et al. 2003). The private forest owners in Sweden, who withdrew from the FSC process, opted for the Programme for Endorsement of Forest Certification (PEFC), which has been successfully in bringing a significant numbers of small private non-industrial forest owners into the certification movement." Some part of the forestry industry may perceive FSC standards as being incompatible with “plantation forestry”. Importantly, a "planted forest" is not necessarily a "plantation" (as defined in FSC standard⁸⁶) since it may have most of the principle characteristics and key elements of native forest ecosystems indigenous to an area.

standards are specifically designed for plantation management (others for natural forest management or NTFP). In 2014/15 (after years of multi-stakeholder negotiations) the SC Std 01-001Version 5 was endorsed and it came with FSC STD 60-004 – the International Generic Indicators (IGI). So far, and probably until Oct 2015, no forest will be certified against Version 5 – all certification is based on Version 4. In P&C Std 01-001 V5 there is not any longer the “P1-9 plus P10 plantation”: “In terms of vegetation, the P&C are globally applicable to all types and scales of forest including natural forests, plantations (...)”. FSC STD 60-004 – distinguish only in few cases (e.g. requirements for regeneration) between plantations and other forest management types.

⁸² The Pan-European Certification Scheme, which is supported by private forest owner associations, was launched in 1999 as a response to the FSC and was later renamed Programme for the Endorsement of Forest Certification Schemes [www.pefc.org] (Gulbrandsen 2005; Johansson & Lidestav 2011).

⁸³ FSC 2012. FSC Principles and Criteria for Forest Stewardship. Document reference code: FSC-STD-01-001 V5-0 EN. Approval date: 10 February 2012. Forest Stewardship Council. <https://ic.fsc.org/download.fsc-std-01-001-v5-0-revised-principles-and-criteria-for-forest-stewardship.a-1780.pdf> [Accessed May 2014].

⁸⁴ The person or entity holding or applying for certification and therefore responsible for demonstrating compliance with the requirements upon which FSC certification is based (Source: FSC 2011).

⁸⁵ A species, subspecies or lower taxon, introduced outside its natural past or present distribution; includes any part, gametes, seeds, eggs, or propagules of such species that might survive and subsequently reproduce (Source: Convention on Biological Diversity (CBD), Invasive Alien Species Programme. Glossary of Terms as provided on CBD website) Source: FSC 2012.

⁸⁶ Plantation: A forest area established by planting or sowing with using either alien or native species, often with one or few species, regular spacing and even ages, and which lacks most of the principal characteristics and key elements of natural forests. Forest Stewardship Council (2015) FSC International Standard. FSC Principles and Criteria for Forest Stewardship, FSC-STD-01-001 V5-2 EN (<https://ic.fsc.org/en/certification/principles-and-criteria/the-revised-pc>). One of the ten FSC principles (Principle 10) addresses plantations directly (<https://ic.fsc.org/en/certification/principles-and-criteria/the-10-principles>).

Most certification standards refer to the use of appropriate provenances, varieties and species for afforestation and reforestation. Native species are always preferred, but alien species are allowed where they are substantially superior to indigenous species for reaching plantation objectives (Stupak et al. 2011)⁸⁷.

The international FSC standard states that native species are preferred, but alien tree species are tolerated as long as their use is monitored and carefully controlled, and adverse ecological effects are avoided (Criterion 6.9). Native species are also preferred in plantations, but aliens can be used when they perform better (Criterion 10.4). In the context of FSC a genetically modified organism is defined as an “organism in which the genetic material has been altered in a way that does not occur naturally by mating and/or natural recombination or both”. The use of GM trees is generally prohibited (Criterion 6.8).

FSC Papua New Guinea⁸⁸ prohibits field tests of GMOs, and some national standards give further details related to alien species and require, for example, a system to be place to monitor spontaneous regeneration outside plantation areas, unusual mortality, disease, insect outbreaks or other adverse environmental impacts (SW Australia), that alien species are only permitted in stands as single trees or in small groups (FSC Luxembourg⁸⁹), or that they only be used in plantations or plant nurseries, or if needed to maintain historical places (FSC Russia)⁹⁰ (Stupak et al. 2011).

⁸⁷ Cf. also FSC STD 01-001 V4: 6.2 Safeguards shall exist which protect rare, threatened and endangered species and their habitats (...). 6.3 Ecological functions and values shall be maintained intact, enhanced, or restored, including: (a) Forest regeneration and succession; (b) Genetic, species, and ecosystem diversity; (c) Natural cycles that affect the productivity of the forest ecosystem. 6.9 The use of exotic species shall be carefully controlled and actively monitored to avoid adverse ecological impacts. 10.2 The design and layout of plantations should promote the protection, restoration and conservation of natural forests, and not increase pressures on natural forests. Wildlife corridors, (...). 10.4 The selection of species for planting shall be based on their overall suitability for the site and their appropriateness to the management objectives. In order to enhance the conservation of biological diversity, native species are preferred over exotic species in the establishment of plantations and the restoration of degraded ecosystems. Exotic species, which shall be used only when their performance is greater than that of native species, shall be carefully monitored to detect unusual mortality, disease, or insect outbreaks and adverse ecological impacts.

⁸⁸ (<https://ic.fsc.org/papua-new-guinea.285.htm>). This is supposed to be the same for all FSC National Standards, as the P&C say: FSC STD 01-001 V4:8 Use of biological control agents (...); Use of genetically modified organisms shall be prohibited. FSC STD 01-001 V5: 10.4 The Organization shall not use genetically modified organisms in the Management Unit. Field testing within the scope of the certificate is therefore also not permitted – some national indicators stress this, others not. But the result is the same: No GMO in any FSC certified FMUs

⁸⁹ <http://www.fsc-lux.lu/>

⁹⁰ Luxemburg and Russia FSC are examples where the national standard requires more than the global P&C. Similarly, e.g., FSC in Germany, where they say no plantations, except for Christmas trees, and not larger than 5 ha resp. 5% of the forest management unit. FSC Germany: 6.9.1 Tree species that are not part of natural forest associations (including exotic species) are positioned as single trees or small groups to an extent which does not jeopardize the long-term development of the stands into natural forest associations. 6.9.1.1 If the proportion of tree species that are not part of natural forest associations exceeds 20% of the planned stocking goal for the specific forest management unit, the forest enterprise professionally justifies that the development towards the natural forest plant association is not at risk. 6.9.1.2 Such proof is not necessary for nurse crop that is not part of natural forest associations, if at most 20% of the stocking unit is taken over as temporary mixture. 6.9.1 Tree species that are not part of natural forest associations (including exotic species) are positioned as single trees or small groups to an extent which does not jeopardize the long-term development of the stands into natural forest associations..9.2 Positioning of tree species that are not part of natural forest associations (including exotic species) in areas that fall under principle 9, is only feasible insofar as it is explicitly permitted by the respective environmental sector planning (e.g. protective area regulation, Natura 2000 management plan). For the Christmas tree planted forests it says: 10.4.3 Exotic species are attentively monitored to avoid negative impacts on the forest ecosystem. The forest enterprise makes sure that negative impacts are avoided through the use of appropriate measures. Additionally, the addenda to FSC Germany, at 6.9.1.1 states that non-native tree species are only cultivated in Germany when they have been proven ecologically non-invasive through years of experience or with comparable data from pilot projects. That is, they must coexist with native tree species and not tend toward dominance. They must support an abundant level of plant and animal life that

Native species are also generally preferred by PEFC (see the Pan European Operational Level Guidelines, PEOLG⁹¹), but the use of alien species is allowed as long as negative impacts can be avoided or minimized.

4.5 European initiatives and legislation

4.5.1 *Habitat Directive – Natura 2000*

The Habitats Directive⁹² (together with the Birds Directive) forms the cornerstone of Europe's nature conservation policy. It is built around two pillars: the Natura 2000 network of protected sites and the strict system of species protection. All in all, the directive protects over 1,200 animals and plant species and over 200 habitat types (e.g. special types of forests, meadows, wetlands, etc.), which are of European interest. According to Article 22.b, in implementing the provisions of this Directive, Member States shall: “ensure that the deliberate introduction into the wild of any species which is not native to their territory is regulated so as not to prejudice natural habitats within their natural range or the wild native fauna and flora and, if they consider it necessary, prohibit such introduction. The results of the assessment undertaken shall be forwarded to the committee for information”.

BOX 4.5.1.1 - Management of Natura 2000 habitats: the 9360 Macaronesian laurel forests (Laurus, Ocotea).

Widespread throughout mainland Europe before the Ice Ages (during the Neogene), the humid to hyper-humid evergreen forests known as laurel forests were driven close to extinction during cold climatic periods. Now restricted to the cloud belt of the Macaronesian islands, they grow in deep soils at between 500 and 1,500 m. Macaronesian laurel forests have been intensively transformed since the fifteenth century when the original forest area was largely razed to create farmland and degraded due to forest exploitation and livestock farming. Already significantly reduced in extent, in some areas habitat is being further degraded by exploitation and livestock stocking. In some cases, habitat reduction has led to fragmentation, threatening habitat diversity and leading to species extinction. Other current threats are the spread of alien species, a major concern in the Azores and Madeira, and forest fires, especially serious in the Canary Islands. Native forests have been cleared for pastures but have also been replanted since 1940's with planted forests of alien tree species such as *Acacia* spp. and *Cryptomeria japonica* (Hervías Parejo et al. 2014).

Where necessary, the type of management is chosen according to the degree of habitat development and to local features. The most common situations are: selective cuttings to improve regeneration in stands that have been heavily exploited, conversion of planted forests into laurel forests, control of invasive alien species and recovery of specific threatened species.

In the case of Tenerife (Canary Islands), Arevalo et al. (2011) suggest that conservation and restoration efforts have to be devoted to removal of current *E. globulus* plantations and gradual thinning of *P. radiata*, with the final objective of converting the current planted forests to forests resembling the laurel forests in structure and composition.

is not significantly under those of natural forest plant associations. They must contribute to the performance of the forest's ecological function and regenerate naturally under existing environmental conditions (<https://ic.fsc.org/germany.278.htm>).

⁹¹ (http://pefc.org/images/documents/MCPFE_PEOLG.pdf).

⁹² Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora. Official Journal L 206, 22/07/1992 P. 0007 – 0050 [http://ec.europa.eu/environment/nature/legislation/habitatsdirective/index_en.htm].

Management of Natura 2000 habitats⁹³ is a project launched by the European Commission in January 2007 aimed at defining best practices for management of habitat types included in Annex I of the Habitat Directive (92/43/EEC) that need active recurring management. Twenty-six habitat types that are representative of different bio-geographical regions have been considered. This scenario motivated several LIFE European financed programs aimed at laurel forest restoration.

4.5.2 The Plant Health Regime in the European Union

European Union rules on plant health⁹⁴ aim to protect crops, fruit, vegetables, flowers, ornamentals and forests from harmful pests and diseases (harmful organisms) by preventing their introduction into the EU or their spread within the EU. This aim helps to contribute to the protection of public and private green spaces, forests and the natural landscape. Although the main focus is the control of harmful organisms (pests) within the Community, as a result, the introductions of some tree species might be restricted or specifically regulated due to phytosanitary reasons⁹⁵.

4.5.3 The Biodiversity Strategy of the European Union

In 2011, the European Commission adopted a new strategy that lays down the framework for EU action over the next ten years in order to meet the 2020 biodiversity headline target set by EU leaders in 2010⁹⁶.

The Target 5 of the EU Biodiversity strategy requires that “by 2020 Invasive Alien Species (IAS) and their pathways are identified and prioritised, priority species are controlled or eradicated, and pathways are managed to prevent the introduction and establishment of new IAS”. Within the Action 16 of the Target 5 the EU has committed itself a dedicated legislative instrument on the issue.

⁹³ http://ec.europa.eu/environment/nature/natura2000/management/best_practice_en.htm

⁹⁴ EU rules on plant health form the EU Plant Health Regime which the Commission has reviewed for the first time since 1977. The Commission has proposed a new EU plant health regulation in May 2013. Council Directive 2000/29/EC (Council Directive 2000/29/EC of 8 May 2000, on protective measures against the introduction into the Community of organisms harmful to plants or plant products and against their spread within the Community) provides the basis for this aim. The general principles are based upon provisions laid down in the International Plant Protection Convention (IPPC). Directive 2000/29/EC is supported by a number of Control Directives and Emergency Measures. Cf also Commission Implementing Directive (EU) 2017/1279 of 14 July 2017 amending Annexes I to V to Council Directive 2000/29/EC on protective measures against the introduction into the Community of organisms harmful to plants or plant products and against their spread within the Community.

⁹⁵ The Commission Implementing Decision of 1 March 2012 as regards emergency measures to prevent the introduction into and the spread within the Union of *Anoplophora chinensis* (Forster) (notified under document C(2012) 1310) (2012/138/EU - Official Journal of the European Union L 64/38, of 3 March 2012) banned the introduction of that plants of *Acer* spp. into the Union until 30 April 2012. Under EU Plant Health Regime some tree species are prohibited to be imported from non-EU countries (as listed in Annex 111), e.g. imports of *Chamaecyparis* spp. are banned from countries outside of the EU.

⁹⁶ <http://ec.europa.eu/environment/nature/info/pubs/docs/factsheets/Biod%20Strategy%20FS.pdf>

4.5.4 *The EU Regulation on invasive alien species*

A Regulation on invasive alien species has been adopted by the European Parliament and by the Council on the 22 October 2014⁹⁷ and came into force on 1 January 2015. This legislation seeks to address the problem of invasive alien species in a comprehensive manner so as to protect native biodiversity and ecosystem services, as well as to minimize and mitigate the human health or economic impacts that these species can have⁹⁸.

The core of Regulation (EU) 1143/2014 is a list of invasive alien species of Union concern (Union list). This list is developed on the basis of scientifically underpinned risk assessments, with the assistance of Member States.

The legislation foresees three types of interventions to address species on the Union list; prevention, early detection and rapid eradication, and management. Species on the Union list may not be intentionally brought into the territory of the Union, nor may they be kept, bred, transported to, from or within the Union, placed on the market, grown or released into the environment. Member States need to address their pathways of unintentional introduction and spread. The Regulation also requires that Member States establish a surveillance system for early detection and apply measures for rapid eradication of newly establishing species or management measures for widely spread species. Furthermore, Member States must provide for penalties if the regulation is not correctly applied.

The initial list of invasive alien species of Union concern entered into force on 3 August 2016. The first update of the Union list entered into force on 2 August 2017⁹⁹. As a result, the Consolidated List of Invasive Alien Species of Union concern¹⁰⁰ includes a total of 49 species, with a total number of 23 plant species. At the moment, the Regulation does not address any alien tree species, but only a woody shrub (*Baccharis halimifolia*).

4.5.5 *Forest policies in the European Union*

Forest policies in the European Union are implemented by Member States within a clearly defined framework of established ownership rights and with a long history of national and regional laws and regulations based on long term planning. Although the Treaties for the European Union make no provision for a common forest policy, there is a long history of EU measures supporting certain forest-related activities, coordinated with Member States mainly through the Standing Forestry Committee (European Commission 2003).

Forests are affected by a broad array of Community policies and initiatives arising from diverse EU sectoral policies (e.g., Schmithüsen et al. 2000). For several decades now, environmental forest functions have attracted increasing attention mainly in relation to the protection of biodiversity and, more recently, in the context of climate change impacts and policies. In public perception, apart from the traditional production of wood and other forest products, forests are increasingly valued for their role as public amenities, biodiversity reservoirs, regulators of climate and local weather, sources of clean water, protection against natural disasters and renewable energy sources¹⁰¹ (European Commission 2003).

⁹⁷ Regulation (EU) No 1143/2014 of the European Parliament and of the Council of 22 October 2014 on the prevention and management of the introduction and spread of invasive alien species, published in the Official Journal of the European Union, L 317, 4.11.2014, p. 35–55 (<http://eur-lex.europa.eu/legal-content/EN/TXT/?qid=1415726405933&uri=CELEX:32014R1143>).

⁹⁸ http://ec.europa.eu/environment/nature/invasivealien/index_en.htm

⁹⁹ Commission Implementing Regulation (EU) 2017/1263 of 12 July 2017 updating the list of invasive alien species of Union concern established by Implementing Regulation (EU) 2016/1141 pursuant to Regulation (EU) No 1143/2014 of the European Parliament and of the Council.

¹⁰⁰ http://ec.europa.eu/environment/nature/invasivealien/list/index_en.htm

¹⁰¹ <http://ec.europa.eu/environment/forests/fpolicies.htm>

In 1992, the European Commission launched a programme to increase afforestation activities on farmland¹⁰². The purpose of the program was to reduce the costs of agricultural subsidies. Landowners willing to convert agricultural land into forest or woodland received afforestation grants which included a cost support for maintenance during the first critical years as well as forest premium compensation for the income lost from agricultural products (Dohrenbusch & Bolte 2007). Within the first decade of the programme's launch, about a million ha were afforested in the European Community, mainly in Spain, Portugal, and Ireland. Countries implementing this program were allowed some flexibility, within a limited framework, to modify tree species (including non-native trees), grants and premiums permitted¹⁰³ (Dohrenbusch & Bolte 2007; Lefebvre et al. 2012).

The EU Forestry Strategy¹⁰⁴, first adopted in 1998, put forward as its overall principles the application of sustainable forest management and the multifunctional role of forests. In line with the principle of subsidiarity, meaning that every administrative decision should always been made at the most appropriate level taking into account the specific local circumstances, this Strategy seeks to establish a coherent framework of forest-related actions at EU level. It also aims to improve the linkages and co-ordination between different policy areas as well as the coherence with the forest policies of the Member States (European Commission 2003).

The contents of the Council Directive 1999/105/EC of 22 December 1999 on the marketing of forest reproductive material are also noteworthy¹⁰⁵.

¹⁰² Council Regulation No 2080/1992 of 30 June 1992(OJ L215, 30.7.1992). See also the Reg.No. 2078/1992.

¹⁰³ In Ireland, for example, afforestation grants differed between 2,000 Euro and 5,000 Euro per hectare, dependent on tree species composition. Plantations of conifers such as Sitka spruce (*Picea sitchensis*), for example, or lodgepole pine (*Pinus contorta*), with some 2,500 plants per hectare, attracted a grant of about 2,000 Euro plus 700 Euro for maintenance. For broadleaved species, such as the common oak (*Quercus robur*) or the European beech (*Fagus sylvatica*), the afforestation grant was more than 5,000 Euro along with maintenance compensation of 1,600 Euro. A forest premium is paid up to 20 years for farmers but only 15 years for non-farmers (Dohrenbusch & Bolte 2007).

¹⁰⁴ Council Resolution of 15 December 1998 on a forestry strategy for the European Union. (OJ C56, 26.2.1999).

¹⁰⁵ [Official Journal L 011, 15/01/2000 P. 0017 - 0040]. This Directive contains specific definitions that – although solely referring to the purposes of the Directive itself - are somewhat different from the generally agreed definitions of alien and native status (Cf. Article 2). "... (d) Autochthonous and indigenous means either of the following: (i) Autochthonous stand or seed source: An autochthonous stand or seed source is one which normally has been continuously regenerated by natural regeneration. The stand or seed source may be regenerated artificially from reproductive material collected in the same stand or seed source or autochthonous stands or seed sources within the close proximity; (ii) Indigenous stand or seed source: An indigenous stand or seed source is an autochthonous stand or seed source or is a stand or seed source raised artificially from seed, the origin of which is situated in the same region of provenance. (e) Origin: For an autochthonous stand or seed source, the origin is the place in which the trees are growing. For a non-autochthonous stand or seed source, the origin is the place from which the seed or plants were originally introduced. The origin of a stand or seed source may be unknown. (f) Provenance: The place in which any stand of trees is growing. (g) Region of Provenance: For a species or sub-species, the region of provenance is the area or group of areas subject to sufficiently uniform ecological conditions in which stands or seed sources showing similar phenotypic or genetic characters are found, taking into account altitudinal boundaries where appropriate ...".

In 2006 the EU underpinned its support for sustainable forest management and the multifunctional role of forests by adopting an EU Forest Action Plan¹⁰⁶. The plan was a framework for forest-related measures and was used to coordinate EU initiatives with the forest policies of the Member States. There were 18 key actions proposed to be implemented jointly with the Member States during the period 2007-2011.

In 2014, the EU adopted a revised Forest Strategy¹⁰⁷ which responds to the new challenges facing forests and the forest sector. In the Strategy and in the accompanying documentation, it is stressed that European forests are threatened by biotic and abiotic agents, such as insects and other pests, diseases, grazing and invasive alien species, windstorms, forest fires, droughts, floods and avalanches. Importantly, this Strategy does not list the forest sector as a potential pathway and driver for the introduction and dissemination of new invasive alien species. In 2014 The Council adopted conclusions which welcome the new EU Forest Strategy, underlining the need to enhance forests' adaptive capacities and resilience to climate change, to reduce the risks and effects of forest fires, pests and diseases and invasive alien species and other disturbances with preventive measures.

Rural development policy is part of the EU's common agricultural policy (CAP) which has been the main instrument for implementing forestry measures in recent years. In this context, financial support from the EU for forestry measures, not including direct funding by the Member States, amounted to EUR 4 800 million for the period 2000–2006 (almost 10 % of the rural development budget, source: EC 2011 – Eurostat).

High Nature Value (HNV) forestry can be defined as natural forests and those semi-natural forests in Europe where the management (historical or present) supports a high diversity of native species and habitats and/or which support the presence of species of European, and/or national, and/or regional conservation concern (European Commission 2013). The maintenance and enhancement of HNV farming and forestry systems is a strategic objective of the European Rural Development Policy and the Managing Authority has to monitor and assess the effectiveness of rural development measures as regards this objective. In order to perform the assessment, the European Commission has envisaged three indicators for HNV farmlands and forestry, in the context of the Common Monitoring and Evaluation Framework (CMEF) for Rural Development 2007-2013 (see EC Reg. no. 1974/2006): baseline indicator 18, result indicator 6 and impact indicator 5. The application of these indicators is a challenging task, mainly due to the complexity of the concept to be measured (Pignatti et al. 2012).

4.5.6 The EU Energy Policy

Bio-energy is seen as one of the key options to mitigate greenhouse gas emissions and substitute fossil fuels (e.g., Faaij 2006). As a result, the large-scale production of renewable heat, electricity and transport fuel from biomass is an important component in many climate change mitigation and energy supply scenarios and a strategically important option for increasing the global uptake of renewable energy (Slade et al. 2014). Yet the practicalities of accelerating deployment are mired in controversy over the potential resource conflicts that might occur, particularly over land, water, biodiversity conservation, soil fertility (Slade et al. 2014; Somerville et al. 2010) and forest conservation (Biello 2011). This calls into question whether policies to promote bioenergy are always and everywhere justified (Slade et al. 2014; Searle & Malins 2014) and if they could constitute a further pathway for the introduction of new alien trees.

¹⁰⁶ The Action Plan does not refer directly to the risk posed by invasive alien species. Anyway, the Key action 7 is titled: Contribute towards achieving the revised Community biodiversity objectives for 2010 and beyond. It can be considered as a commitment to the principles aiming to tackle invasive alien species that are present in the recalled document, i.e. the Commission Communication of 22 May 2006 "Halting the loss of biodiversity by 2010 - and beyond - Sustaining ecosystem services for human well-being" [COM(2006) 216 final - Not published in the Official Journal].

¹⁰⁷ Brussels, 20.9.2013, COM(2013) 659 final - http://ec.europa.eu/agriculture/forest/strategy/index_en.htm

At both the international level and the EU level there are several ongoing initiatives seeking more sustainable forest management also in relation to bioenergy from forests and planted forests, such as the various international processes on SFM based on C&I, reducing emissions from deforestation and forest degradation (REDD+) or forest law enforcement, governance and trade (FLEGT). At the European level, the revision of the EU Forestry Strategy (COM/2013/0659 “EU Forest Strategy: for the forests and the forest-based sector”) and the ongoing process to establish a legally binding agreement on forests in Europe must be underlined. In addition to those general provisions for forests and forestry, several specific initiatives for woody bioenergy have been developed with the aim of guaranteeing overall environmental sustainability (Fritsche et al. 2014; Repo et al. 2015)¹⁰⁸.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge all the colleagues and experts that have provided useful information on invasive alien trees, national black lists, forestry management issues and on other parts of this Code of Conduct, and in particular, Paulina Anastasiu, Trausti Baldursson, Linda Berglund, Urszula Biereźnoj-Bazille, Etienne Branquart, Sarah Brunel, Wim Buysse, Ignazio Camarda, Thomas Campagnaro, Paulo Carmo, María Amparo Carrillo-Gavilán, Alberto Del Lungo, Pierre Ehret, René Eschen, Franz Essl, Astra Garkaje, Quentin Groom, Richard Howe, Melanie Josefsson, Marion Karmann, Walter Kollert, Frank Krumm, Sandrine Liegeois, Merike Linnamagi, Cristina Máguas, Elizabete Marchante, Hélia Marchante, Mariam Mironova, Andrei Orlinski, Gerardo Sánchez Peña, Ewa Pisarczyk, Peter Roberntz, Helen Roy, Joaquim Sande Silva, Lisa Schembri, Tommaso Sitzia, Wojciech Solarz, Øystein Størkersen, Rob Tanner, Teodora Trichkova, Margarida Tomé, Maurits Vandegehuchte, Lucie Vitkova, Vladimir Vladimirov, Gian-Reto Walther and Pawel Wasowicz.

DMR acknowledges funding from the DST-NRF Centre of Excellence for Invasion Biology in South Africa and the National Research Foundation of South Africa (grant 85417).

5. REFERENCES

- Aarrestad PA, Myking T, Stabbetorp OE, Tollefsrud MM (2014) Foreign Norway spruce (*Picea abies*) provenances in Norway and effects on biodiversity. NINA Report 1075, 39 pp.
- Adriaens T, Sutton-Croft M, Owen K, Brosens D, van Valkenburg J, Kilbey D, Groom Q, Ehmig C, Thürkow F, Van Hende P, Schneider K (2015) Trying to engage the crowd in recording invasive alien species in Europe: experiences from two smartphone applications in northwest Europe. *Management of Biological Invasions* 6(2): 215-225.
- Aitken SN, Bemmels JB (2016) Time to get moving: assisted gene flow of forest trees. *Evolutionary Applications*, 9: 271–290.
- Aitken SN, Yeaman S, Holliday JA, Wang T, Curtis-McLane S (2008) Adaptation, migration or extirpation: climate change outcomes for tree populations. *Evolutionary Applications* 1: 95-111.
- Águas A, Ferreira A, Maia P et al. (2014) Natural establishment of *Eucalyptus globulus* Labill. in burnt stands in Portugal. *For Ecol Manage* 323: 47-56.
- Aguilar S, Condit R (2001) Use of native tree species by an Hispanic community in Panama. *Economic Botany* 55: 223-235.
- Aguilera J, Nielsen KM, Sweet J (2013) Risk assessment of GM trees in the EU: current regulatory framework and guidance. *iForest* 6: 127-131.
- Allen J (1998) Mangroves as alien species: the case of Hawaii. *Global Ecology & Biogeography Letters* 7: 61-71.

¹⁰⁸ Cf. Also the BASIS project on Biomass Availability and Sustainability Information System (http://www.basisbioenergy.eu/fileadmin/BASIS/D4.1.Sustainability_Criteria_for_Bioenergy.pdf).

- Alston KP, Richardson DM (2006) The roles of habitat features, disturbance, and distance from putative source populations in structuring alien plant invasions at the urban/wildland interface on the Cape Peninsula, South Africa. *Biological Conservation* 132: 183-198.
- Anastasiu P, Negrean G (2005) Invasive and potential invasive alien plants in Romania (Black List). In: *Bioplatform – Romanian National Platform for Biodiversity. Inter-Institutional Protocol for Biodiversity Research Development* (ed. Mihăilescu Simona). București: Edit. Academiei Române, pp. 107-114.
- Andersson B, Engelmark O, Rosvall O, Sjöberg K (1999) Miljökonsekvensbeskrivning (MKB) av skogsbruk med contortatall i Sverige. *Skog Forsk Redogörelse* 1, 50 pp. (in Swedish).
- Andreu J, Vilà M, Hulme PE (2009) An assessment of stakeholder perceptions and management of noxious alien plants in Spain. *Environmental Management* 43:1244-1255.
- Arévalo JR, Fernández-Palacios JM (2005) Gradient analysis of exotic *Pinus radiata* plantations and potential restoration of natural vegetation in Tenerife, Canary Islands (Spain). *Acta Oecologica* 27(1): 1-8.
- Arevalo JR, Delgado JD, Fernandez-Palacios JM (2011) Regeneration of potential laurel forest under a native canopy and an exotic canopy, Tenerife (Canary Islands). *Forest Systems* 20(2): 255-265.
- Auld G, Gulbrandsen LH, McDermott CL (2008) Certification. *Annual Review of Environment and Resources* 33: 187-211.
- Backlund I, Bergsten U (2012) Biomass production of dense direct-seeded Lodgepole Pine (*Pinus contorta*) at Short Rotation Periods. *Silva Fennica* 46: 609-623.
- Tanya Baycheva-Mergera T, Wolfslehner B (2016) Evaluating the implementation of the Pan-European Criteria and indicators for sustainable forest management – A SWOT analysis. *Ecological Indicators* 60: 1192–1199.
- Bernier P, Schoene D (2009) Adapting forests and their management to climate change: an overview. *Unasylva* 60: 3-4.
- Bianco P, Ciccacese L, Jacomini C, Pellegrino P (2014) Impacts of short rotation forestry plantations on environments and landscape in Mediterranean basin. *Rapporti* 196/14. ISPRA – Istituto Superiore per la Protezione e la Ricerca Ambientale, Roma, 115 pp.
- Biello D (2011) The false promise of biofuels. *Scientific American* 305: 58-65. Published online: 19 July 2011 | doi:10.1038/scientificamerican0811-58.
- Blackburn TM, Pyšek P, Bacher S, Carlton JT, Duncan RP, Jarošík V, Wilson JR, Richardson DM (2011) A proposed unified framework for biological invasions. *Trends in Ecology & Evolution* 26: 333-339.
- Boerjan W (2005) Biotechnology and the domestication of forest trees. *Current Opinion in Biotechnology* 16: 159-166.
- Bonfils A-C (2006) Canada's regulatory approach. In: Williams CG (ed.), *Landscapes, genomics and transgenic conifers*, Springer, Dordrecht, The Netherlands, pp. 229-243.
- Bonneh O (2000) Management of planted pine forest in Israel: past, present and future. In: Ne'eman G, Trabaud L (eds.) *Ecology, biogeography and management of Pinus halepensis and P. brutia forest ecosystems in the Mediterranean Basin*. Backhuys Publishers, Leiden, The Netherlands, pp. 377-390.
- Boström M (2003) How state-dependent is a non-state driven rule-making project? The case of forest certification in Sweden. *Journal of Environmental Policy and Planning* 5(2): 165-180.
- Boström M (2003) How state-dependent is a non-state driven rule-making project? The case of forest certification in Sweden. *Journal of Environmental Policy and Planning* 5 (2): 165-180.

- Boyd IL, Freer-Smith PH, Gilligan CA, Godfray HCJ (2013) The Consequence of Tree Pests and Diseases for Ecosystem Services. *Science* 342: 823-829.
- Branco M, Brockerhoff EG, Castagneyrol B (2014) Host range expansion of native insects to exotic trees increases with area of introduction and presence of congeneric native trees. *Journal of Applied Ecology*. Doi 10.1111/1365-2664.12362
- Branquart E (ed.) (2014) Alert, black and watch lists of invasive species in Belgium. Harmonia version 1.2, Belgian Forum on Invasive species, accessed on the 10th November 2014 from: <http://ias.biodiversity.be>.
- Bremer LL, Farley KA (2010) Does plantation forestry restore biodiversity or create green deserts? A synthesis of the effects of land-use transitions on plant species richness. *Biodiversity and Conservation* 19: 3893-3915.
- Brockerhoff EG, Jactel H, Parrotta JA, Quine CP, Sayer J (2008) Plantation forests and biodiversity: oxymoron or opportunity? *Biodiversity and Conservation* 17: 925-951.
- Broncano MJ, Vilà M, Boada M (2005) Evidence of *Pseudotsuga menziesii* naturalization in montane Mediterranean forests. *Forest Ecology and Management* 211: 257-263.
- Brooks ML, D'Antonio CM, Richardson DM, Grace JB, Keeley JE, Di Tomaso JM, Hobbs RJ, Pellant M, Pyke D (2004) Effects of invasive alien plants on fire regimes. *BioScience* 54: 677-688.
- Brown NR, Noss RF, Diamond DD, Myers MN (2001) Conservation biology and forest certification working together toward ecological sustainability. *Journal of Forestry* 88: 18-25.
- Brown ND, Curtis T, Adams EC (2015) Effects of clear-felling versus gradual removal of conifer trees on the survival of understorey plants during the restoration of ancient woodlands. *Forest Ecology and Management* 348: 15-22.
- Brundu G (2008) La legislazione italiana in materia di specie vegetali alloctone. In: Galasso G, Chiozzi G, Azuma M, Banfi E (eds.), *Le specie alloctone in Italia: censimenti, invasività e piani di azione*. Memorie della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano 36(1): 44.
- Brunel S, Petter F, Fernandez-Galiano E, Smith I (2009) Approach of the European and Mediterranean Plant Protection Organization to the Evaluation and Management of Risks Presented by Invasive Alien Plants. In: Inderjit (ed.), *Management of Invasive Weeds, Invading Nature - Springer Series in Invasion Ecology*, pp. 319-343.
- Brunner AM, Li J, Di Fazio SP, Shevchenko O, Montgomery BE, Mohamed R, Wei H, Ma C, Elias AA, Van Wormer K, Strauss SH (2007) Genetic containment of forest plantations. *Tree Genetics & Genomes*. doi 10.1007/s11295-006-0067-8.
- Buddenhagen CE, Chimera C, Clifford P (2009) Assessing biofuel crop invasiveness: a case study. *PLoS One* 4, doi: 10.1371/journal.pone.0005261.
- Calviño-Cancela M (2013) Effectiveness of eucalypt plantations as a surrogate habitat for birds. *Forest Ecology and Management* 310: 692-699.
- Calviño-Cancela M, Rubido-Bará M (2013) Invasive potential of *Eucalyptus globulus*: Seed dispersal, seedling recruitment and survival in habitats surrounding plantations. *Forest Ecology and Management* 305: 129-137.
- Canadell G, Raupach MR (2008) Managing Forests for Climate Change Mitigation. *Science* 320: 1456-1457.
- Capdevila-Argüelles L, Zillett B (2008) A perspective on climate change and invasive alien species. T-PVS/Inf (2008) 5 rev. 2nd Meeting of the Group of Experts on Biodiversity and Climate Change.

- Convention on the Conservation of European Wildlife and Natural Habitats, Standing Committee, Strasbourg, 16 June 2008, 31 pp.
- Caplat P, Hui C, Maxwell BD, Peltzer DA (2014) Cross-scale management strategies for optimal control of trees invading from source plantations. *Biological Invasions* 16: 677-690.
- Carnus J-M, Parrotta J, Brockerhoff E, Arbez M, Jactel H, Kremer A, Lamb D, O'Hara K, Walters B (2006) Planted Forests and Biodiversity. *Journal of Forestry* 104(2): 65-77.
- Carrillo-Gavilán MA, Vilà M (2010) Little evidence of invasion by alien conifers in Europe. *Diversity and Distributions* 16: 203-213.
- Cashore B, Auld G, Newsom D (2004) *Governing Through Markets - Forest Certification and the Emergence of Non-State Authority*. Yale University Press, New Haven.
- Castro-Díez P, Valle G, González-Muñoz N, Alonso A (2014) Can the life-history strategy explain the success of the exotic trees *Ailanthus altissima* and *Robinia pseudoacacia* in Iberian floodplain forests? *PloS One* 9(6), doi: 10.1371/journal.pone.0100254.
- Catry FX, Moreira F, Deus E, Silva JS, Águas A (2015) Assessing the extent and the environmental drivers of *Eucalyptus globulus* wildling establishment in Portugal: results from a countrywide survey. *Biological Invasions*, DOI 10.1007/s10530-015-0943-y.
- Cierjacks A, Kowarik I, Joshi J, Hempel S, Ristow M, Lippe M, Weber E (2013) Biological flora of the British Isles: *Robinia pseudoacacia*. *Journal of Ecology* 101(6): 1623-1640.
- Chimera CC, Buddenhagen CE, Clifford PM (2010) Biofuels: the risks and dangers of introducing invasive species. *Biofuels* 1: 785-796.
- Christen B, Dalgaard T (2012) Buffers for biomass production in temperate European agriculture: A review and synthesis on function, ecosystem services and implementation, *Biomass and Bioenergy*. doi: 10.1016/j.biombioe.2012.09.053.
- Ciccarese L, Pellegrino P, Silli V, Zanchi G (2014) Short rotation forestry and methods for carbon accounting. A case study of black locust (*Robinia pseudoacacia* L.) plantation in central Italy. *Rapporti 200/2014*, ISPRA - Istituto Superiore per la Protezione e la Ricerca Ambientale, Roma, 49 pp.
- Council of Europe (1977) *Liste des plantes rares, menacées et endémiques en Europe*. Collection Sauvegarde de la Nature. Strasbourg, France.
- Corlett RT (2005) Vegetation. In: Gupta A (ed), *The Physical Geography of Southeast Asia*. Oxford University Press, Oxford, pp. 105-119.
- Courbet F, Lagacherie M, Marty P, Ladier J, Ripert C, Riou-Nivert P, Huard F, Amandier L, Paillassa É (2012) *Le cèdre en France face au changement climatique: bilan et recommandations*.
- Craven D, Hall JS, Verjans JM (2008) Impacts of herbicide application and mechanical cleanings on growth and mortality of two timber species in *S. spontaneum* grasslands of the Panama Canal watershed. *Restoration Ecology* doi:10.1111/j.1526-100x.2008.00408.x
- DAISIE. 2009. *Handbook of alien species in Europe*. Knoxville, TN: Springer.
- D'Antonio C, Meyerson LA (2002) Exotic plant species as problems and solutions in ecological restoration: a synthesis. *Restoration Ecology* 10: 703-713.
- Daehler CC, Denslow JS, Ansari S, Kuo H-C (2004) A risk assessment system for screening out invasive pest plants from Hawai'i and other Pacific Islands. *Conservation Biology* 18: 360-368.
- Danin A (2000) The inclusion of adventive plants in the second edition of *Flora Palaestina*. *Willdenowia* 30: 305-314.

- Dauber J, Jones MB, Stout JC (2010) The impact of biomass crop cultivation on temperate biodiversity. *GCB Bioenergy* 2: 289-309.
- Declan L, Collins K, Cross J, Cooke D, McGinnity P (2014) Native Riparian Woodlands - A Guide to Identification, Design, Establishment and Management. Native Woodland Scheme Information Note No. 4 [<http://www.woodlandsofireland.com/sites/default/files/Riparian.pdf>].
- den Herder M, Burgess PJ, Mosquera-Losada MR, Herzog F, Hartel T, Upton M, Viholainen I, Rosati A (2015) Preliminary stratification and quantification of agroforestry in Europe. Milestone Report 1.1 for EU FP7 Research Project: AGFORWARD 613520. (22 April 2015). 57 pp [<http://www.agforward.eu/index.php/en/preliminary-stratification-and-quantification-of-agroforestry-in-europe.html>].
- Deniz T, Yildirim HT (2014) Institutional and political assessment of forest plantations in Turkey. *International Forestry Review* 16(2): 199-204.
- Despain DG (2001) Dispersal ecology of lodgepole pine (*Pinus contorta* Dougl.) in its native environment as related to Swedish forestry. *Forest Ecology and Management* 141: 59-68.
- Di Fazio SP, Leonardi S, Slavov GT, Garman SL, Adams WT, Strauss SH (2012) Gene flow and simulation of transgene dispersal from hybrid poplar plantations. *New Phytologist* 193: 903-915.
- Dia A, Duponnois R (2010) Le projet majeur africain de la Grande Muraille Verte - Concepts et mise en œuvre. IRD, Institut de Recherche pour le Développement, Marseille, France, 439 pp.
- Dias FS, Bugalho MN, Cerdeira JO, Martins MJ (2013) *Biodiversity and Conservation* 22: 93-112.
- Diaz S, Hector A, Wardle DA (2009) Biodiversity in forest carbon sequestration initiatives: not just a side benefit, *Current Opinion in Environmental Sustainability* 1(1): 55-60.
- Dickie IA, Bennett BM, Burrows LE, Nuñez MA, Peltzer DA, Porté A, Richardson DM, Rejmánek M, Rundel PW, van Wilgen BW (2014) Conflicting values: ecosystem services and invasive tree management. *Biological Invasions* 16: 705-719.
- Dodet M, Collet C (2012) When should exotic forest plantation tree species be considered as an invasive threat and how should we treat them? *Biological Invasions* 14: 1765-1778.
- Donaldson JE, Hui C, Richardson DM, Wilson JRU, Robertson MP, Webber BL (2014) Invasion trajectory of alien trees: the role of introduction pathway and planting history. *Global Change Biology* 20: 1527-1537.
- Dohrenbusch A, Bolte A (2007) Forest Plantations. In: Kües U (ed.), *Wood Production, Wood Technology, and Biotechnological Impacts*, Universitätsverlag Göttingen, Germany, pp. 73-84.
- Ducci (2014) Species restoration through dynamic ex situ conservation: *Abies nebrodensis* as a model. In: Bozzano M, Jalonen R, Thomas E, Boshier D, Gallo L, Cavers S, Bordács S, Smith P, Loo J (eds.), *The state of the world's forest genetic resources – Thematic study. Genetic considerations in ecosystem restoration using native tree species*, pp. 225-233, FAO, Rome.
- Drinan TJ, Graham CT, O'Halloran J, Harrison SSC (2013) The impact of conifer plantation forestry on the Chydoridae (Cladocera) communities of peatland lakes. *Hydrobiologia* 700: 203-219.
- Dye P, Caren C (2004) Water use by black wattle (*Acacia mearnsii*): implications for the link between removal of invading trees and catchment streamflow response. *South African Journal of Science* 100: 40-44.
- Eggleton M, Zegada-Lizarazu W, Ephrath J, Berliner P (2007) The effect of brackish water irrigation on the above- and below-ground development of pollarded *Acacia saligna* shrubs in an arid environment. *Plant Soil* 299: 141-152.

- Eichhorn MP, Paris P, Herzog F, Incoll LD, Liagre F, Mantzanas K, Mayus M, Moreno G, Papanastasis VP, Pilbeam DJ, Pisanelli A, Dupraz C (2006). Silvoarable systems in Europe - past, present and future prospects. *Agroforestry Systems* 67: 29-50.
- Elfving B, Norgren O (1993) Volume yield superiority of lodepole pine compared to Scots pine in Sweden. In: *Pinus contorta - from untamed forest to domesticated crop*. Department of Genetics and Plant Physiology, Swedish University of Agricultural Sciences, Umeå, Report 11, pp 69-80.
- Elwes HJ, Henry AH (1910) *The Trees of Great Britain and Ireland*. S.R. Publishers, Wakefield (reprint 1971).
- Engelmark O, Sjöberg K, Andersson B, Rosvall O, Ågrend GO, Baker WL, Barklund P, Björkman C, Don G, Despainf, Elfving B, Ennos RA, Karlman M, Knecht MF, Knight DH, Ledgard NJ, Lindelöw Å, Nilsson C, Peterken GF, Sörlin S, Sykes MT (2001) Ecological effects and management aspects of an exotic tree species: the case of lodgepole pine in Sweden. *Forest Ecology and Management* 141: 3-13.
- EPPO (2009) PP 1/141(3): Weeds in tree and shrub nurseries. *EPPO Bulletin* 39(3): 246-249.
- Essl F (2005): Verbreitung, Status und Habitatbindung der subspontanen Bestände der Douglasie (*Pseudotsuga menziesii*) in Österreich. *Phyton* 45/1: 117-144.
- Essl F, Dullinger S, Rabitsch W, Hulme PE, Hülber K, Jarošík V, Kleinbauer I, Krausmann F, Kühn I, Nentwig W, Vilà M, Genovesi P, Gherardi F, Desprez-Loustau M-L, Roques A, Pyšek P (2011) Socioeconomic legacy yields an invasion debt. *Proceedings of the National Academy of Sciences of the United States of America* 108(1): 203-207.
- Essl F, Nehring S, Klingenstein F, Milasowszky N, Nowack C, Rabitsch W (2011) Review of risk assessment systems of IAS in Europe and introducing the German-Austrian Black List Information System (GABLIS). *Journal for Nature Conservation* 19: 339-350.
- Etienne M (2000) Pine agroforestry in the West Mediterranean basin. In: Ne'eman G, Trabaud L (eds) *Ecology, biogeography and management of Pinus halepensis and P. brutia forest ecosystems in the Mediterranean Basin*. Backhuys Publishers, Leiden, The Netherlands, pp. 355-368.
- European Commission (2003) *Sustainable forestry and the European Union - Initiatives of the European Commission*. Luxembourg: Office for Official Publications of the European Communities, 56 pp.
- European Commission (2013) *Communication from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions. A new EU Forest Strategy: for forests and the forest-based sector*. Brussels, 20.9.2013.
- European Commission (2016) *Sustainable forest management* (http://ec.europa.eu/growth/sectors/raw-materials/industries/forest-based/sustainable-forest-management/index_en.htm).
- European Environment Agency (2008) *European forests - ecosystem conditions and sustainable use*. EEA Report number 3/2008, EEA, Copenhagen.
- Evans J (1980) Prospects for eucalypts as forest trees in Great Britain. *Forestry* 53(2): 129-43.
- Evans J (2009a) The Multiple Roles of Planted Forests. In: Evans J (ed) *Planted forests: uses, impacts, and sustainability*. CAB International and FAO, pp. 61-90.
- Evans J (2009b) Sustainable Silviculture and Management. In: Evans J (ed.) *Planted forests: uses, impacts, and sustainability*. CAB International and FAO, pp. 113-140.
- Faaij APC (2006) Bio-energy in Europe: changing technology choices. *Energy Policy*: 322-342.
- FAO (2001) *Global Forest Resources Assessment 2000, Main Report*, FAO Forestry Paper 140, Food and Agriculture Organization of the United Nations, Rome, Italy, 479 pp. [<ftp://ftp.fao.org/docrep/fao/003/Y1997E/FRA%202000%20Main%20report.pdf>].

- FAO (2006a). Global planted forests thematic study. Results and analysis. Planted Forests and Trees Working Paper No. FP38. Food and Agriculture Organization of the United Nations, Food and Agriculture Organization of the United Nations, Rome, Italy.
- FAO (2006b) Responsible management of planted forests: voluntary guidelines. Planted Forests and Trees Working Paper 37/E, Food and Agriculture Organization of the United Nations, Rome, Italy, 73 pp. [www.fao.org/forestry/site/10368/en].
- FAO (2006c) Fire management: voluntary guidelines. Principles and strategic actions. Fire Management Working Paper 17. Food and Agriculture Organization of the United Nations, Rome, Italy, 63 pp. [www.fao.org/forestry/site/35853/en].
- FAO (2010a) Global Forest Resources Assessment 2010. Main Report. FAO Forestry Paper 163. Food and Agriculture Organization of the United Nations, Rome, Italy, 340 pp.
- FAO (2010b) Fighting sand encroachment - Lessons from Mauritania. FAO Forestry Paper 158, Food and Agriculture Organization of the United Nations, Rome, Italy, 74 pp.
- FAO (2010c) Planted forests in sustainable forest management. A statement of principles. Food and Agriculture Organization of the United Nations, Rome, Italy, 16 pp. [<http://www.fao.org/docrep/012/al248e/al248e00.pdf>].
- FAO (2010d) Forests and genetically modified trees. Food and Agriculture Organization of the United Nations, Rome, Italy, 235 pp. [<http://www.fao.org/docrep/013/i1699e/i1699e.pdf>].
- FAO (2011) Guide to implementation of phytosanitary standards in forestry. FAO Forestry Paper 164, Food and Agriculture Organization of the United Nations, Rome, Italy, 101 pp.
- FAO (2012) FRA 2015 Terms and Definitions. Forest Resources Assessment Working Paper 180, Food and Agriculture Organization of the United Nations, Rome, Italy, 31 pp. [<http://www.fao.org/docrep/017/ap862e/ap862e00.pdf>].
- FAO (2013) Advancing Agroforestry on the Policy Agenda: A guide for decision-makers, by G. Buttoud, in collaboration with O. Ajayi, G. Detlefsen, F. Place & E. Torquebiau. Agroforestry Working Paper no. 1. Food and Agriculture Organization of the United Nations. FAO, Rome, Italy, 37 pp.
- FAO (2014) Great Green Wall for the Sahara and the Sahel initiative - The African Wall. Food and Agriculture Organization of the United Nations, 6 pp. [<http://www.fao.org/docrep/016/ap603e/ap603e.pdf>].
- FAO (2015a) Global Forest Resources Assessment 2015. Desk reference. Food and Agriculture Organization of the United Nations, Rome, Italy, 244 pp.
- FAO (2015b) Global Forest Resources Assessment 2015. How are the world's forests changing? Food and Agriculture Organization of the United Nations, Rome, Italy, 47 pp.
- Faulkner KT, Robertson MP, Rouget M, Wilson JR (2014) A simple, rapid methodology for developing invasive species watch lists. *Biological Conservation* 179: 25-32.
- Fedorkov A (2010) Variation in shoot elongation patterns in *Pinus contorta* and *Pinus sylvestris* in north-west Russia. *Scandinavian Journal of Forest Research* 25(3): 208-212.
- Felton A, Boberg J, Björkman C, Widenfalk O (2013) Identifying and managing the ecological risks of using introduced tree species in Sweden's production forestry. *Forest Ecology and Management* 307: 165-177.
- Finch O-D, Szumelda A (2007) Introduction of Douglas fir (*Pseudotsuga menziesii* (Mirb.) Franco) into Western Europe: Epigeic arthropods in intermediate-aged pure stands in northwestern Germany. *Forest Ecology and Management* 242: 260-272.

- Finnoff D, Shogren J, Leung B, Lodge D (2007) Take a risk: preferring prevention over control biological invaders. *Ecological Economics* 62: 216-222.
- Fischer A, Vasseur L (2002) Smallholder perceptions of agroforestry projects in Panama. *Agrofor Syst* 54: 103-113. doi:10.1023/A:1015047404867
- FITEC (2007) Practice for Plantation Forestry. Best Environmental Practices. New Zealand, 1st Edition, 57 pp.
- FOREST EUROPE, UNECE, FAO (2011) State of Europe's Forests 2011. Status and Trends in Sustainable Forest Management in Europe. Ministerial Conference on the Protection of Forests in Europe, Forest Europe Liaison Unit Oslo, 337 pp.
- FOREST EUROPE (2016) What is Forest Europe? (http://www.foresteuropa.org/about_us/foresteuropa).
- Forestry Commission (2011) Forests and Climate Change. UK Forestry Standard Guidelines. Forestry Commission, Edinburgh. i-iv + 1-68 pp.
- Fossati T, Grassi F, Sala F, Castiglione S (2003) Molecular analysis of natural populations of *Populus nigra* L. intermingled with cultivated hybrids. *Molecular Ecology* 12: 2033-2043.
- Framstad E (ed) (2009) Increased biomass harvesting for bioenergy - effects on biodiversity, landscape amenities and cultural heritage values. *TemaNord* 2009:591, Nordic Council of Ministers, Copenhagen [www.norden.org/publications].
- Frankenhuyzen K von, Beardmore T (2004) Current status and environmental impact of transgenic trees. *Canadian Journal of Forest Research* 34: 1163-1180.
- Friedman J, Auble G, Shafroth P, Scott M, Merigliano M, Freehling M, Griffin E (2005) Dominance of non-native riparian trees in western USA. *Biological Invasions* 7: 747-751.
- Fritsche UR, Iriarte L, de Jong J, Agostini A, Scarlat N (2014) Extending the EU Renewable Energy Directive sustainability criteria to solid bioenergy from forests. *Natural Resources Forum* 38: 129-140.
- Froude VA (2011) Wilding Conifers in New Zealand: Beyond the status report. Report prepared for the Ministry of Agriculture and Forestry. Pacific Eco-Logic, Bay of Islands. 44p.
- Gaertner M, Biggs R, Te Beest M, Hui C, Molofsky J, Richardson DM (2014) Invasive plants as drivers of regime shifts: Identifying high priority invaders that alter feedback relationships. *Diversity and Distributions* 20: 733-744.
- Gaertner M, Richardson DM, Privett SDJ (2011) Effects of alien plants on ecosystem structure and functioning and implications for restoration: insights from three degraded sites in South African fynbos. *Environmental Management* 48: 57-69.
- García-de-Lomas J, Vilà M (2015) Lists of harmful alien organisms: Are the national regulations adapted to the global world? *Biological Invasions* 17 (11): 3081-3091.
- Garen EJ, Saltonstall K, Slusser JL, Mathias S, Ashton MS, Hall JS (2009) An evaluation of farmers' experiences planting native trees in rural Panama: implications for reforestation with native species in agricultural landscapes. *Agroforestry Systems* doi: 10.1007/s10457-009-9203-4
- Gederaas L, Moen TL, Skjelseth S, Larsen L-K (eds.) (2012) Alien species in Norway - with the Norwegian Black List 2012. The Norwegian Biodiversity Information Centre, Norway, 212 pp.
- Gellini R, Grossoni P (1996) *Botanica Forestale – I – Gimnosperme*. Casa Editrice Antonio Milani, Milano, Italy, 267 pp.

- Genovesi P, Shine C (2004) European strategy on invasive alien species. Convention on the Conservation of European Wildlife and Habitats (Bern Convention). Nature and environment, No. 137, Council of Europe Publishing, Strasbourg, 67 pp.
- Gill AM (1997) Eucalypts and fires: interdependent or independent? In: Williams JE, Woinarski JCZ (eds.), *Eucalypt Ecology: Individuals to Ecosystems*. Cambridge University Press, Cambridge, pp. 151-167.
- Goes E (1991) *A fl oresta portuguesa, sua importância e descrição das espécies de maior interesse*. Edição Portucel, Lisboa, 257 pp.
- Gomez-Zamalloa MG, Caparros A, Ayanz AS-miguel (2011) 15 years of forest certification in the European Union. Are we doing things right? *Forest Systems* 20: 81-94.
- González-García S, Gasol CM, Moreira MT, Gabarrell X, Rieradevall i Pons J, Feijoo G (2011) Environmental assessment of black locust (*Robinia pseudoacacia* L.)-based ethanol as potential transport fuel. *The International Journal of Life Cycle Assessment* 16: 465-477.
- González-Muñoz N, Costa-Tenorio M, Espigares T (2011) Invasion of alien *Acacia dealbata* on Spanish *Quercus robur* forests: Impact on soils and vegetation. *Forest Ecology and Management* 269: 214-221.
- González-Muñoz N, Castro-Diez P, Parker IM (2012) Differences in nitrogen use strategies between native and exotic tree species: predicting impacts on invaded ecosystems. *Plant and Soil* 363: 319-329.
- González-Muñoz N, Linares JC, Castro-Díez C, Sass-Klaassen U (2014) Predicting climate change impacts on native and invasive tree species using radial growth and twenty-first century climate scenarios. *European Journal of Forest Research*. doi: 10.1007/s10342-014-0823-5
- Goosem M, Harding EK, Chester G, Tucker N, Harriss C, Oakley K (2010) *Roads in Rainforest: Best Practice Guidelines for Planning, Design and Management*. Guidelines prepared for the Queensland Department of Transport and Main Roads and the Australian Government's Marine and Tropical Sciences Research Facility. Published by the Reef and Rainforest Research Centre Limited, Cairns, 64 pp.
- Gordon DR, Tancig KJ, Onderdonk DA, Gantz CA (2011) Assessing the invasive potential of biofuel species proposed for Florida and the United States using the Australian Weed Risk Assessment. *Biomass Bioenerg* 35: 74-79.
- Gordon DR, Flory SL, Cooper AL, Morris SK (2012) Assessing the invasion risk of *Eucalyptus* in the United States using the Australian Weed Risk Assessment. *International Journal of Forestry Research*, doi:10.1155/2012/203768.
- Grattapaglia D, Kirst M (2008) *Eucalyptus* applied genomics: from gene sequences to breeding tools. *New Phytologist* 179: 911-929.
- Gray LK, Gylander T, Mbogga MS, Chen P-Y, Hamann A (2011) Assisted migration to address climate change: Recommendations for aspen reforestation in western Canada. *Ecological Applications* 21: 1591-1603.
- Grünewald H, Böhm C, Quinkenstein A, Grundmann P, Eberts J, von Wühlisch G (2009) *Robinia pseudoacacia* L.: a lesser known tree species for biomass production. *BioEnergy Research* 2: 123-133.
- Gulbrandsen LH (2005) The effectiveness of non-state governance schemes: a comparative study of forest certification in Norway and Sweden. *International Environmental Agreements* 5: 125-149.

- Gumiero B, Mant J, Hein T, Elso J, Boz B (2013) Linking the restoration of rivers and riparian zones/wetlands in Europe: Sharing knowledge through case studies. *Ecological Engineering* 56: 36-50.
- Gunnarsson KS, Eysteinnsson T, Curl SL, Thorfinnsson T (2005) Iceland. *Acta Silv. Lign. Hung. Special Edition*: 335-346 [http://aslh.nymh.hu/fileadmin/dokumentumok/fmk/acta_silvatica/cikkek/VolE1-2005/iceland.pdf].
- Hadincová V, Köhnleinová I, Marešová J (2007) Invasive behaviour of white pine (*Pinus strobus*) in sandstone areas in the Czech Republic. In: Härtel H, Cílek V, Herben T, Jackson A, Williams R, editors. *Sandstones Landscapes*. Praha: Academia. 219-224.
- Häggman H, Raybould A, Borem A, Fox T, Handley L, Hertzberg M, Lu M-Z, Macdonald P, Oguchi T, Pasquali G, Pearson L, Peter G, Quemada H, Séguin A, Tattersall K, Ulian E, Walter C, McLean M (2013) Genetically engineered trees for plantation forests: key considerations for environmental risk assessment. *Plant Biotechnology Journal*. doi: 10.1111/pbi.12100
- Halford M, Heemers L, van Wesemael D, Mathys C, Wallens S, Branquart E, Vanderhoeven S, Monty A, Mahy G (2014) The voluntary Code of conduct on invasive alien plants in Belgium: results and lessons learned from the AlterIAS LIFE+ project. *EPPO Bulletin* 44(2): 212-222.
- Hamrick JL (1990) Isozymes and the analysis of genetic structure in plant populations. *Isozymes in Plant Biology* (eds D.E. Soltis & P.S. Soltis), pp. 87–105. Chapman & Hall, London, UK.
- Haysom KA, Murphy ST (2003) The status of invasiveness of forest tree species outside their natural habitat: a global review and discussion paper. *Forest Health and Biosecurity Working Paper FBS/3E*. Forestry Department. FAO, Rome (unpublished) [<http://www.fao.org/docrep/006/J1583E/J1583E00.HTM>].
- Hardcastle PD (2006) A review of the impacts of short-rotation forestry. LTS International. *Short Rotation Coppice Willow, Best Practice Guidelines* (Ireland).
- Harwood CE, Nambiar EKS (2014) Productivity of Acacia and Eucalypt plantations in Southeast Asia. 2. Trends and variations. *International Forestry Review* 16(2): 249-260.
- Hawkins CL, Bacher S, Essl F, Hulme PE, Jeschke JM, Kühn I, Kumschick S, Nentwig W, Pergl J, Pyšek P, Rabitsch W, Richardson DM, Vilà M, Wilson JRU, Genovesi P, Blackburn TM (2015) Framework and guidelines for implementing the proposed IUCN Environmental Impact Classification for Alien Taxa (EICAT). *Diversity and Distributions* 21: 1360–1363.
- Heleno RH, Ceia RS, Ramos JA, Memmott J (2009) Effects of alien plants on insect abundance and biomass: a food-web approach. *Conservation Biology* 23: 410-419.
- Heller NE, Zavaleta ES (2009) Biodiversity management in the face of climate change: A review of 22 years of recommendations. *Biological Conservation* 142(1): 14-32.
- Hemström K, Mahapatra K, Gustavsson L (2014) Public Perceptions and acceptance of intensive forestry in Sweden. *Ambio* 43(2): 196-206.
- Heritage Council (1999) *Policy Paper on Forestry and the National Heritage*. The Heritage Council, Kilkenny.
- Hermý M, Honnay O, Firbank L, Grashof-Bokdam C, Lawesson JE (1999) An ecological comparison between ancient and other forest plant species of Europe, and the implications for conservation. *Biological Conservation* 91: 9-22.
- Hermý M, Verheyen K (2007) Legacies of the past in the present-day forest biodiversity: a review of past land-use effects on forest plant species composition and diversity. *Ecological Research* 22: 361-371.

- Hervías Parejo S, Ceia RS, Ramos JA, Sampaio HL, Heleno RH (2014) Tiptoeing between restoration and invasion: seed rain into natural gaps within a highly invaded relic forest in the Azores. *European Journal of Forest Research* 133: 383-390.
- Heywood VH, Brunel S (2009) Code of Conduct on Horticulture and Invasive Alien Plants. Nature and Environment No. 155. Strasbourg, Council of Europe Publishing.
- Heywood VH, Brunel S (2011) Code of Conduct on Horticulture and Invasive Alien Plants. Illustrated version. Nature and Environment No. 162. Strasbourg, Council of Europe Publishing.
- Heywood VH, Sharrock S (2013) European Code of Conduct for Botanic Gardens on Invasive Alien Species. Council of Europe, Strasbourg, Botanic Gardens Conservation International, Richmond, 61 pp.
- Hickie D (1997) Evaluation of Environmental Designations in Ireland. The Heritage Council, Kilkenny.
- Higgins SI, Richardson DM (1999) Predicting plant migration rates in a changing world: the role of long-distance dispersal. *American Naturalist* 153: 464-475.
- Hinchee M, Rottmann W, Mullinax L, Zhang C, Chang S, Cunningham M, Pearson L, Nehra N (2009) Short-rotation woody crops for bioenergy and biofuels applications. *In Vitro Cellular & Developmental Biology - Plant* 45: 619-629.
- Hinchee M, Zhang C, Chang S, Cunningham M, Hammond W, Nehra N (2011) Biotech Eucalyptus can sustainably address society's need for wood: the example of freeze tolerant Eucalyptus in the southeastern US. *BMC Proc* 5(Suppl. 7): 124.
- Hobbs RJ, Higgs E, Harris JA (2009) Novel ecosystems: implications for conservation and restoration. *Trends in Ecology and Evolution* 24: 599-605.
- Hoenicka H, Fladung M (2006) Biosafety in *Populus* spp. and other forest trees: from non-native species to taxa derived from traditional breeding and genetic engineering. *Trees - Structure and Function* 20: 131-144.
- Hughes FMR (ed.) (2003) *The Flooded Forest: Guidance for policy makers and river managers in Europe on the restoration of floodplain forests*. FLOBAR2, Department of Geography, University of Cambridge, UK. 96 pp.
- Iverson LR, Prasad AM, Matthews SN, Peters M (2008) Estimating potential habitat for 134 eastern US tree species under six climate scenarios. *Forest Ecology and Management* 254: 390-406. doi: 10.1016/j.foreco.2007.07.023
- Ivors K, Garbelotto M, Vries IDE, Ruyter-Spira C, Hekkert BTE, Rosenzweig N, Bonants P (2006) Microsatellite markers identify three lineages of *Phytophthora ramorum* in US nurseries, yet single lineages in US forest and European nursery populations. *Molecular Ecology* 15: 1493-1505.
- Jackson RB, Jobbágy EG, Avissar R, Baidya Roy S, Barrett DJ, Cook CW, Farley KA, le Maitre DC, McCarl BA, Murray BC (2005) Trading Water for Carbon with Biological Carbon Sequestration. *Science* 310: 1944-1947.
- Jama B, Zeila A (2005) *Agroforestry in the drylands of eastern Africa: a call to action*. ICRAF Working Paper no. 1. Nairobi: World Agroforestry Centre, 29 pp.
- James RR, Di Fazio SP, Brunner AM, Strauss SH (1998) Environmental effects of genetically engineered woody biomass crops. *Biomass & Bioenergy* 14: 403-414.
- Janse G. Ed. (2008) *Best Practices in Forest Communication. Contributions from the Forest Communicators Network*. UNECE-FAO Forest Communicators Network, 61 pp.
- Jansson S, Douglas CJ (2007) *Populus: A Model System for Plant Biology*. *Annual Review of Plant Biology* 58: 435-458.

- Jeschke JM, Bacher S, Blackburn TM, Dick JTA, Essl F, Evans T, Gaertner M, Hulme PE, Kühn I, Mrugała A, Pergl J, Pyšek P, Rabitsch W, Ricciardi A, Richardson DM, Sendek A, Vilà M, Winter M, Kumschick S (2014) Defining the impact of non-native species. *Conservation Biology* 28: 1188-1194.
- Jeschke JM, Keesing F, Ostfeld RS (2013) Novel organisms: comparing invasive species, GMOs, and emerging pathogens. *AMBIO* 42: 541-548.
- Johansson J, Lidestav G (2011) Can voluntary standards regulate forestry? Assessing the environmental impacts of forest certification in Sweden. *Forest Policy and Economics* 13: 191-198.
- Kaonga LM (2010) Fractal analysis of canopy architectures of *Acacia angustissima*, *Gliricidia sepium*, and *Leucaena collinsii* for estimation of aboveground biomass in a short rotation forest in eastern Zambia. *Journal of Forestry Research* 23(1): 1-12.
- Karlman M (2001) Risks associated with the introduction of *Pinus contorta* in northern Sweden with respect to pathogens. *Forest Ecology and Management* 141: 97-105.
- Karp A, Shield I (2008) Bioenergy from plants and the sustainable yield challenge. *New Phytologist* 179: 15-32.
- Katsanevakis S, Genovesi P, Gaiji S, Nyegaard Hvid H, Roy H, Nunes AL, Sánchez Aguado F, Bogucarskis K, Debusscher B, Deriu I, Harrower C, Josefsson M, Lucy FE, Marchini A, Richards G, Trichkova T, Vanderhoeven S, Zenetos A, Cardoso AC (2013) Implementing the European policies for alien species – networking, science, and partnership in a complex environment. *Management of Biological Invasions* 4: 3–6.
- Kawaletz H, Mölder I, Zerbe S, Annighöfer P, Terwei A, Ammer C (2013) Exotic tree seedlings are much more competitive than natives but show underyielding when growing together. *Journal of Plant Ecology*: 1-11, doi: 10.1093/jpe/rts044.
- Kellezi M, Stafasani M, Kortoci Y (2012) Evaluation of biomass supply chain from *Robinia pseudoacacia* L. SRF plantations on abandoned lands. *Journal of Life Sciences* 6: 187-193.
- Kim TN, Matsumura J, Oda K (2011) Effect of growing site on the fundamental wood properties of natural hybrid clones of *Acacia* in Vietnam. *Journal of Wood Science* 57: 87-93.
- Kjær ED, Lobo A, Myking T (2014) The role of exotic tree species in Nordic forestry. *Scandinavian Journal of Forest Research* 29(4): 323–332.
- Kleinbauer I, Dullinger S, Peterseil J, Essl F (2010) Climate change might drive the invasive tree *Robinia pseudoacacia* into nature reserves and endangered habitats. *Biological Conservation* 143: 382-390.
- Knapic S, Pirralho M, Louzada JL, Pereira H (2014) Early assessment of density features for 19 *Eucalyptus* species using X-ray microdensitometry in a perspective of potential biomass production. *Wood Science and Technology* 48: 37-49.
- Knapic S, Tavares F, Pereira H (2006) Heartwood and sapwood variation in *Acacia melanoxylon* R. Br. Trees in Portugal. *Forestry* 79 (4), doi:10.1093/forestry/cpl010.
- Korosuo A, Sandström P, Öhman K, Eriksson LO (2014) Impacts of different forest management scenarios on forestry and reindeer husbandry. *Scandinavian Journal of Forest Research* 29 (1): 234-251.
- Kraus D, Krumm F (eds.) (2013) Integrative approaches as an opportunity for the conservation of forest biodiversity. European Forest Institute. 284 pp.
- Kreyling J, Bittner T, Jaeschke A, Jentsch A, Steinbauer MJ, Thiel D, Beierkuhnlein C (2011) Assisted colonization: A question of focal units and recipient localities. *Restoration Ecology* 19: 433-440.

- Krivánek M, Pyšek P (2006) Predicting invasions by woody species in a temperate zone: a test of three risk assessment schemes in the Czech Republic (Central Europe) *Diversity and Distributions* 12: 319-327.
- Kuehne C, Nosko P, Horwath T, Bauhus J (2014) A comparative study of physiological and morphological seedling traits associated with shade tolerance in introduced red oak (*Quercus rubra*) and native hardwood tree species in southwestern Germany. *Tree Physiology* 00, 1-10. doi:10.1093/treephys/tpt124
- Kull CA, Shackleton CM, Cunningham PJ, Ducatillon C, Dufour-Dror J-M, Esler K.J, Friday JB, Gouveia AC, Griffin AR, Marchante E, Midgley SJ, Pauchard A, Rangan H, Richardson DM, Rinaudo T, Tassin J, Urgenson LS, von Maltitz GP, Zenni RD, Zylstra MJ (2011) Adoption, use and perception of Australian acacias around the world. *Diversity and Distributions* 17: 822-836.
- Kumschick S, Richardson DM (2013) Species-based risk assessments for biological invasions: advances and challenges. *Diversity and Distributions* 19: 1095-1105.
- Kuzovkina YA, Weih M, Romero MA, Charles J, Hurst S, McIvor I, Karp A, Trybush S, Labrecque M, Teodorescu I et al. (2008) *Salix*: botany and global horticulture. *Horticultural Reviews* 34: 448-490.
- Lamb D, Erskine P, Parrotta JA (2005) Restoration of degraded tropical forest landscapes. *Science* 310:1628-1632. doi:10.1126/science.1111773
- Lavi A, Perevolotsky A, Kigel J, Noy-Meir I (2005) Invasion of *Pinus halepensis* from plantations into adjacent natural habitats. *Applied Vegetation Science* 8: 85-92.
- Ledford H (2014) Brazil considers transgenic trees. *Nature* 512: 357,
- Lefebvre M, Espinosa M, Gomez y Paloma S (2012) The influence of the Common Agricultural Policy on agricultural landscapes. Report EUR 25459 EN. Luxembourg: Publications Office of the European Union, 78 pp.
- Le Maitre DC, Gaertner M, Marchante E, Ens E-J, Holmes PM, Pauchard A, O'Farrell PJ, Rogers AM, Blanchard R, Blignaut J, Richardson DM (2011) Impacts of invasive Australian acacias: implications for management and restoration. *Diversity and Distributions* 17: 1015-1029.
- Le Maitre DC, van Wilgen BW, Gelderblom CM, Bailey C, Chapman RA, Nel JA (2002) Invasive alien trees and water resources in South Africa: case studies of the costs and benefits of management. *Forest Ecology and Management* 160: 143-159.
- Le Maitre DC, Versfeld DB, Chapman RA (2000) The impact of invading alien plants on surface water resources in South Africa: a preliminary assessment. *Water SA* 26, 397-408.
- Ledgard N (2002) The spread of Douglas-fir into native forests. *New Zealand Journal of Forestry* 47: 36-38.
- Leite A, Santos C, Saraiva I, Pinho JR (1999) O planeamento florestal e as espécies invasoras. In: 1º Encontro de Invasoras Lenhosas. Gerês 16 - 18 Nov. Sociedade Portuguesa de Ciências Florestais, Lisboa, pp. 45-50.
- Leslie AD, Mencuccini M, Perks M (2012) The potential for Eucalyptus as a wood fuel in the UK. *Applied Energy* 89: 176-182.
- Leung B, Roura-Pascual N, Bacher S, Heikkilä J, Brotons L, Burgman MA, Dehnen-Schmutz K, Essl F, Hulme PE, Richardson DM, Sol D, Vilà M (2012) TEASing apart alien species risk assessments: a framework for best practices. *Ecology Letters* 15: 1475-1493.
- Levin E (2005) Growing China's great green wall. *ECOS* 13: 127. [http://www.ecomagazine.com/?act=view_file&file_id=EC127p13.pdf]

- Li MS (2006) Ecological restoration of mineland with particular reference to the metalliferous mine wasteland in China: a review of research and practice. *Science of the Total Environment* 357: 38-53.
- Liebholt AM, Brockerhoff EG, Garrett LJ, Parke JL, Britton KO (2012) Live plant imports: the major pathway for forest insect and pathogen invasions of the US. *Frontiers in Ecology and the Environment* 10(3): 135-143, doi:10.1890/110198.
- Lindenmayer DB, Hulvey KB, Hobbs RJ, Colyvan M, Felton A, Possingham H, Steffen W, Wilson K, Youngtob K, Gibbons P (2012) Avoiding bio-perversity from carbon sequestration solutions. *Conservation Letters* 5: 28-36.
- Liphshitz N, Biger G (2001) Past distribution of Aleppo pine (*Pinus halepensis*) in the mountains of Israel (Palestine). *Holocene* 11: 427-436.
- Locatelli B, Kanninen M, Brockhaus M. et al. (2008) Facing an uncertain future: how forests and people can adapt to climate change. *Forest Perspectives No. 5*. CIFOR, Bogor, Indonesia.
- Loehr RC, Haith DA, Walter MF, Martin CS (1979) *Best Management Practices for Agriculture and Silviculture*. Ann Arbor, MI: Ann Arbor Science Publishers Inc. 740 pp.
- Lopian R (2005) The International Plant Protection Convention and invasive alien species. In: IPPC Secretariat. 2005. Identification of risks and management of invasive alien species using the IPPC framework. Proceedings of the workshop on invasive alien species and the International Plant Protection Convention, Braunschweig, Germany, 22–26 September 2003. FAO Rome, Italy, pp 6-16.
- Lorentz KA, Minogue PJ (2015) Exotic Eucalyptus plantations in the southeastern US: risk assessment, management and policy approaches. *Biological Invasions* 17: 1581-1593.
- Lorenzo P, González L, Reigosa MJ (2010) The genus *Acacia* as invader: the characteristic case of *Acacia dealbata* Link in Europe. *Annals of Forest Science* 67: 101, doi: 10.1051/forest/2009082.
- Lorenzo P, Palomera-Pérez A, Reigosa MJ, González L (2011) Allelopathic interference of invasive *Acacia dealbata* Link on the physiological parameters of native understory species. *Plant Ecology* 212: 403-412.
- Love B, Spaner D (2005) A survey of small-scale farmers using trees in pastures in Herrera Province, Panama. *Journal of Sustainable Forestry* 20:37-65. doi:10.1300/J091v20n03_03.
- Lowe S, Browne M, Boudjelas S (2000) 100 of the world's worst invasive alien species. A selection from the global invasive species database. Invasive Species Specialist Group, Auckland, New Zealand.
- Lugo AE (1997) The apparent paradox of reestablishing species richness on degraded lands with tree monocultures. *Forestry Ecology and Management* 99: 9-19.
- Mabey R (1996) *Flora Britannica*. Sinclair-Stevenson, London.
- McCarthy J, Canziani O, Leary N. et al. (eds) (2001). *Climate change 2001: impacts, adaptation, and vulnerability*. Contribution of Working Group II to the third assessment report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, UK.
- Macdonald IAW, Graber DM, DeBenedetti S, Groves RH, Fuentes ER (1988) Introduced species in nature reserves in Mediterranean-type climatic regions of the world. *Biological Conservation* 44: 37-66.
- Madsen CL, Dahl CM, Thirslund KB, Grousset F, Johannsen VK, Ravn HP (2014) Pathways for non-native species in Denmark. Department of Geosciences and Natural Resource Management, University of Copenhagen, Frederiksberg. 131 pp. [<http://ign.ku.dk/formidling/publikationer/rapporter/filer-2014/pathways-for-non-native-species-in-DK.pdf>].

- Madsen CL, Dahl CM, Thirslund KB, Grousset F, Johannsen VK, Ravn HP (2014) Pathways for non-native species in Denmark. Department of Geosciences and Natural Resource Management, University of Copenhagen, Frederiksberg. 131 pp.
- Malcolm DC, Mason WL, Clarke GC (2001) The transformation of conifer forests in Britain - regeneration, gap size and silvicultural systems. *Forest Ecology and Management* 151: 7-23.
- Maltoni A, Mariotti B, Tani A (2012) La gestione della robinia in Toscana. La gestione dei popolamenti, l'impiego in impianti specializzati, il controllo della diffusione. DEISTAF – Dipartimento di Economia, Ingegneria, Scienze e Tecnologie Agrarie e Forestali, Università di Firenze, Centro stampa Giunta Regione Toscana (In Italian) [<http://www.regione.toscana.it/documents/10180/24010/La+gestione+della+robinia+in+Toscana+-+la+gestione+dei+popolamenti,%20l'impiego+in+impianti+specializzati,%20il+controllo+della+diffusione/023b90d8-1a7a-42c5-b200-648767748863?version=1.0>].
- Mandák B, Hadincová V, Mahelka V, Wildová R (2013) European invasion of North American *Pinus strobus* at large and fine scales: high genetic diversity and fine-scale genetic clustering over time in the adventive range. *PLoS One* 8(7): e68514. doi:10.1371/journal.pone.0068514.
- Marchante E, Kjølner A, Struwe S, Freitas H (2008a) Short- and long-term impacts of *Acacia longifolia* invasion on the belowground processes of a Mediterranean coastal dune ecosystem. *Applied Soil Ecology* 40: 210-217.
- Marchante E, Kjølner A, Struwe S, Freitas H (2008b) Invasive *Acacia longifolia* induce changes in the microbial catabolic diversity of sand dunes. *Soil Biology and Biochemistry* 40: 2563-2568.
- Marchante H (2001) Invasão dos ecossistemas dunares portugueses por *Acacia*: uma ameaça para a biodiversidade nativa. Master Thesis. Faculty of Sciences and Technology, University of Coimbra, Coimbra.
- Marchante H, Marchante E, Freitas H (2003) Invasion of the Portuguese dune ecosystems by the exotic species *Acacia longifolia* (Andrews) Willd.: effects at the community level. In: Child LE, Brock JH, Brundu G, Prach K, Pyšek P, Wade PM, Williamson M (eds), *Plant Invasion: Ecological Threats and Management Solutions*. Backhuys Publishers, Leiden, The Netherlands, pp. 75-85.
- Marchante E, Marchante H, Morais MC, Freitas H (2010) Combining methodologies to increase public awareness about invasive plants in Portugal. In: Brunel S, Uludag A, Fernandez-Galiano E, Brundu G (eds.), *2nd International Workshop on Invasive Plants in Mediterranean Type Regions of the World.*, At Trabzon, Turkey. 2-6 August 2010, pp. 227-239.
- Maringera J, Wohlgemuth T, Neff C, Boris Pezzatti G, Conedera M (2012) Post-fire spread of alien plant species in a mixed broad-leaved forest of the Insubric region. *Flora* 207: 19-29.
- Marshall W (1852) Excessive and noxious increase of *Udora canadensis*. *Phytologist* 4: 705-715.
- Marshall W (1857) The American water-weed *Anacharis alsinastrum*. *Phytologist* 2: 194-197.
- Mascheretti S, Croucher PJP, Vettraino A, Prospero S, Garbelotto M (2008) Reconstruction of the sudden oak death epidemic in California through microsatellite analysis of the pathogen *Phytophthora ramorum*. *Molecular Ecology* 17: 2755-2768.
- Masiero M, Secco L, Pettenella D, Brotto L (2015) Standards and guidelines for forest plantation management: A global comparative study. *Forest Policy and Economics* 53: 29-44.
- Maslin BR, Miller JT, Seigler DS (2003) Overview of the generic status of *Acacia* (Leguminosae: Mimosoideae). *Australian Systematic Botany* 16: 1-18.
- Mazzotti FJ, Center TD, Dray FA, Thayer D (2014) Ecological Consequences of invasion by *Melaleuca quinquenervia* in South Florida Wetlands: Paradise Damaged, not Lost. SSWEC123, Wildlife

Ecology and Conservation Department, UF/IFAS Extension. Original publication date June 1997. Reviewed June 2014, 5 pp. [<http://edis.ifas.ufl.edu/uw123>].

- McCormick N, Howard G (2013) Beating back biofuel crop invasions: guidelines on managing the invasion risk of biofuel developments. *Renew Energy* 49: 263-266.
- McGarrigle M, Clenaghan C (2004) Agriculture and Forestry. In: State of Ireland's Environment 2004. Environmental Protection Agency. EPA, Johnstown Castle Estate, Co. Wexford, pp 130-136.
- McKay H (ed) (2011) Short Rotation Forestry: review of growth and environmental impacts. Forest Research Monograph, 2, Forest Research, Surrey, 212 pp.
- McKenney DW, Pedlar JH, Rood RB, Price D (2011) Revisiting projected shifts in the climate envelopes of North American trees using updated general circulation models. *Global Change Biology* 17: 2720-2730. doi: 10.1111/j.1365-2486.2011.02413.x.
- McNeely JA (ed.) (2001) *The Great Reshuffling: Human Dimensions of Invasive Alien Species*. IUCN, Gland, Switzerland and Cambridge, UK. vi + 242 pp.
- Mead DJ (2013) Sustainable management of *Pinus radiata* plantations. FAO Forestry Paper No. 170. Food and Agriculture Organization of the United Nations, Rome, Italy, 246 pp.
- Medina-Villar S, Castro-Díez P, Alonso A, Cabra-Rivas I, Parker MI, Pérez-Corona E (2015) Do the invasive trees, *Ailanthus altissima* and *Robinia pseudoacacia*, alter litterfall dynamics and soil properties of riparian ecosystems in Central Spain? *Plant Soil*, doi: 10.1007/s11104-015-2592-4.
- Meidinger E (2011) Forest certification and democracy. *European Journal of Forest Research* 130: 407-419.
- Meirmans PG, Lamothe M, Gros-Louis M-C, Khasa D, Périnet P, Bousquet J, Isabel N (2010) Complex patterns of hybridization between exotic and native North American Poplar species. *American Journal of Botany* 97(10): 1688-1697.
- MEPA (2002) Guidelines on Trees, Shrubs and Plants for Planting and Landscaping in the Maltese Islands. Environmental Management Unit Planning Directorate, 63 pp. [<http://www.mepa.org.mt/>].
- Messines J (1952) Sand-dune fixation and afforestation in Libya. *Unasyva* 6(2): [<http://www.fao.org/docrep/x5363e/x5363e02.htm#sand%20dune%20fixation%20and%20afforestation%20in%20libya>].
- Messinger J, Güney A, Zimmermann R, Ganser B, Bachmann M, Remmele S, Aas G (2015) *Cedrus libani*: A promising tree species for Central European forestry facing climate change? *European Journal of Forest Research*. doi: 10.1007/s10342-015-0905-z
- Meyerson LA, Mooney HA (2007) Invasive alien species in an era of globalization. *Frontiers in Ecology and the Environment* 5: 199-208.
- Milad M, Schaich H, Konold W (2013) How is adaptation to climate change reflected in current practice of forest management and conservation? A case study from Germany. *Biodiversity and Conservation* 22: 1181-1202.
- Moreira F, Ferreira A, Abrantes N, Catry F, Fernandes P, Roxo L, Keizer J, Silva J (2013) Occurrence of native and exotic invasive trees in burned pine and eucalypt plantations: implications for post-fire forest conversion. *Ecological Engineering* 58: 296-302.
- Morissette S (2012) Is forest certification compatible with fast-growing plantations? Réseau Ligniculture Québec. Information notice, 4 pp. [http://www.poplar.ca/upload/documents/forestcertif_e.pdf].
- Morris TL, Esler KJ, Barger NN, Jacobs SM, Cramer MD (2011) Ecophysiological traits associated with the competitive ability of invasive Australian acacias. *Diversity and Distributions* 17: 898-910.

- Moss T, Monstadt J (2008) Restoring floodplains in Europe. Policy context and project experiences. IWA Publishing, London, UK, 355 pp.
- Mwangi E, Swallow B (2005) Invasion of *Prosopis juliflora* and local livelihoods: Case study from the lake Baringo area of Kenya. ICRAF Working Paper, no. 3. Nairobi: World Agroforestry Centre, 66 pp.
- Nearby DG (2013), Best management practices for forest bioenergy programs. WIREs Energy Environ 2: 614-632. doi: 10.1002/wene.77
- Nearby DG, Smethurst PJ, Baillie B, Petrone KC (2011) Water Quality, Biodiversity and Codes of Practice in Relation to Harvesting Forest Plantations in Streamside Management Zones. CSIRO, 100 pp.
- Nocentini S (2010) Le specie forestali esotiche: la sperimentazione di Aldo Pavari e le prospettive attuali. L'Italia Forestale e Montana / Italian Journal of Forest and Mountain Environments 65 (4): 449-457.
- Notov AA, Vinogradova YuK, Mayorov SR (2011) On the problem of development and management of regional black books. Russian Journal of Biological Invasions 2(1): 35-45.
- O'Brien EK, Mazanec RA, Krauss SL (2007) Provenance variation of ecologically important traits of forest trees: implications for restoration. Journal of Applied Ecology 44: 583-593.
- Orwig DA (2002) Ecosystem to regional impacts of introduced pests and pathogens: historical context, questions and issues. Journal of Biogeography 29: 1471-1474.
- Osem Y, Lavi A, Rosenfeld A (2011) Colonization of *Pinus halepensis* in Mediterranean habitats: consequences of afforestation, grazing and fire. Biological Invasions 13: 485-498.
- Pairon M, Petitpierre B, Campbell M, Guisan A, Broennimann O, Baret PV, Jacquemart A-L, Besnard G (2010) Multiple introductions boosted genetic diversity in the invasive range of black cherry (*Prunus serotina*; Rosaceae). Annals of Botany 105(6): 881-890.
- Paquette A, Messier C (2010) The role of plantations in managing the world's forests in the Anthropocene. Frontiers in Ecology and the Environment 8(1): 27-34.
- Parker IM, Simberloff D, Lonsdale WM, Goodell K, Wonham M, Kareiva PM, Williamson MH, von Holle B, Moyle PB, Byers JE, Goldwasser L (1999) Impact: Toward a Framework for Understanding the Ecological Effects of Invaders. Biological Invasions 1(1): 3-19.
- Pasta S, Badalamenti E, La Mantia T (2012) *Acacia cyclops* A. Cunn. ex G. Don (Leguminosae) in Italy: first cases of naturalization. Anales del Jardín Botánico de Madrid 69(2): 193-200.
- Pavari A, De Philippis A (1941) La sperimentazione di specie forestali esotiche in Italia. Risultati del primo ventennio. Annali della sperimentazione agraria, vol. XXXVIII, Tip. Failli, Roma, Italy, 646 pp.
- Payn T, Carnus J-M, Freer-Smith P, Kimberley M, Kollert W, Liu S, Orazio C, Rodriguez L, Neves Silva L, Wingfield MJ (2015) Changes in planted forests and future global implications. Forest Ecology and Management 352: 57-67.
- Pedlar JH, McKenney DW, Aubin I, Beardmore T, Beaulieu J, Iverson L, O'Neill GA, Winder RS, Ste-Marie C (2012) Placing forestry in the assisted migration debate. BioScience 62: 835-842.
- Perrings C, Dehnen-Schmutz K, Touza J, Williamson M (2005) How to manage biological invasions under globalization. Trends in Ecology and Evolution 20(5): 212-215.
- Peterken GF (1974) A method for assessing woodland flora for conservation using indicator species. Biological Conservation 6: 39-45.
- Peterken GF (1977) General Management Principles for Nature Conservation in British Woodlands. Forestry 50 (1): 27-48.

- Peterken GF (1981) Woodland conservation and management. Chapman and Hall, London.
- Peterken GF (2001) Ecological effects of introduced tree species in Britain. *Forest Ecology and Management* 141: 31-42.
- Peterken GF, Ausherman D, Buchenau M, Forman RTT (1992) Old-growth Conservation within British Upland Conifer Plantations. *Forestry* 65 (2): 127-144.
- Petitpierre B, Pairon M, Broennimann O, Jacquemart AL, Guisan A, Besnard G (2009) Plastid DNA variation in *Prunus serotina* var. *serotina* (Rosaceae), a North American tree invading Europe. *European Journal of Forest Research* 128: 431–436.
- Pettersson M, Strömberg C, Carina E, Keskitalo H (2016) Possibility to implement invasive species control in Swedish forests. *Ambio* 45(Suppl. 2): S214–S222.
- Pheloung PC, Williams PA, Halloy SR (1999) A weed risk assessment model for use as a biosecurity tool evaluating plant introductions. *Journal of Environmental Management* 57: 239-251.
- Pignatti G, De Natale F, Gasparini P, Mariano A, Trisorio A (2012) High nature value forest areas: a proposal for Italy based on national forest inventory data. *L'Italia Forestale e Montana / Italian Journal of Forest and Mountain Environments* 67(3): 281-288.
- Premoli A, Quiroga P, Gardner M (2013) *Araucaria araucana*. The IUCN Red List of Threatened Species. Version 2014.2. <www.iucnredlist.org>. Downloaded on 21 August 2014.
- Pringle P, Willsman P (2013) Wakatipu Wilding Conifer Control Group & Queenstown Lakes District Council, 33 pp.
- Pyšek P, Prach K (2003) Research into plant invasions in a crossroads region: history and focus. *Biological Invasions* 5: 337-348.
- Pyšek P, Richardson DM, Rejmánek M, Webster G, Williamson M, Kirschner J (2004) Alien plants in checklists and floras: towards better communication between taxonomists and ecologists. *Taxon* 53: 131–143.
- Quine CP, Humphrey JW (2010) Plantations of exotic tree species in Britain: irrelevant for biodiversity or novel habitat for native species? *Biodiversity and Conservation* 19: 1503-1512.
- Radtke A, Ambra S, Zerbe S, Tonon G, Fontana V, Ammer C (2013) Traditional coppice forest management drives the invasion of *Ailanthus altissima* and *Robinia pseudoacacia* into deciduous forests. *Forest Ecology and Management* 291: 308-317.
- Rascher KG, Große-Stoltenberg A, Máguas C, Meira-Neto JAA, Werner C (2011) *Acacia longifolia* invasion impacts vegetation structure and regeneration dynamics in open dunes and pine forests. *Biological Invasions*, doi 10.1007/s10530-011-9949-2.
- Rédei K (2002) Management of black Locust (*Robinia pseudoacacia* L.) stands in Hungary. *Journal of Forestry Research* 13(4): 260-264.
- Rédei K, Csiha I, Keserü Z (2011a) Black locust (*Robinia pseudoacacia* L.) short-rotation crops under marginal site conditions. *Acta Silvatica & Lignaria Hungarica* 7: 125-132.
- Rédei K, Csiha I, Keserü Z, Végh AK, Györi J (2011b) The Silviculture of Black Locust (*Robinia pseudoacacia* L.) in Hungary: a Review. *South-East European Forestry* 2(2): 101-107.
- Reichard SH, Hamilton CW (1997) Predicting invasions of woody plants introduced into North America. *Conservation Biology* 11: 193-203.
- Rejmánek M, Richardson DM (2011) Eucalypts. In: Simberloff D, Rejmánek M (eds), *Encyclopedia of Biological Invasions*, Berkeley and Los Angeles: University of California Press, pp. 203-209.

- Rejmánek M, Richardson DM (2013) Trees and shrubs as invasive alien species - 2013 update of the global database. *Diversity and Distributions* 19: 1093-1094.
- Repo A, Böttcher H, Kindermann G, Liski J (2015) Sustainability of forest bioenergy in Europe: land-use-related carbon dioxide emissions of forest harvest residues. *GCB Bioenergy* 7: 877–887.
- Ribeiro RA, Lemos-Filho JP, Ramos ACS, Lovato MB (2011) Phylogeography of the endangered rosewood *Dalbergia nigra* (Fabaceae): insights into the evolutionary history and conservation of the Brazilian Atlantic Forest. *Heredity* 106: 46-57.
- Ricciardi A, Cohen J (2007) The invasiveness of an introduced species does not predict its impact. *Biological Invasions* 9: 309-315.
- Richardson DM (1998a). Forestry trees as invasive aliens. *Conservation Biology* 12: 18-26.
- Richardson DM (1998b) Invasive alien trees: the price of forestry. In: *Invaders from Planet Earth*. World Conservation Double Issue 4/97-1/98, pp. 14-15. IUCN, Gland, Switzerland.
- Richardson DM, Bond WJ (1991) Determinants of plant distribution: Evidence from pine invasions. *American Naturalist* 137: 639-668.
- Richardson DM, Pyšek P, Rejmánek M, Barbour Mg, Panetta D, West CJ (2000) Naturalization and invasion of alien plants: concepts and definitions. *Diversity and Distributions* 6: 93-107.
- Richardson DM (2011) Forestry and Agroforestry. In: Simberloff D and Rejmánek M (Eds), *Encyclopedia of Biological Invasions*, Berkeley and Los Angeles: University of California Press, pp. 241-248.
- Richardson DM, Binggeli P, Schroth G (2004) Invasive agroforestry trees: problems and solutions. In: Schroth G, de Fonseca GAB, Harvey CA, Gascon C, Vasconcelos HL, Izac A-MN (eds), *Agroforestry and Biodiversity Conservation in Tropical Landscapes*, pp. 371-396. Island Press, Washington, D.C.
- Richardson DM, Carruthers J, Hui C, Impson FAC, Miller JT, Robertson MP, Rouget M, Le Roux JJ, Wilson JRU (2011) Human-mediated introductions of Australian acacias - a global experiment in biogeography. *Diversity and Distributions* 17: 771-787.
- Richardson DM, Holmes PM, Esler KJ, Galatowitsch SM, Stromberg JC, Kirkman SP, Pyšek P, Hobbs RJ (2007) Riparian zones—degradation, alien plant invasions and restoration prospects. *Diversity and Distributions* 13: 126-139.
- Richardson DM, Hellmann JJ, McLachlan JS, Sax DF, Schwartz MW, Gonzalez P, Brennan EJ, Camacho A, Root TL, Sala OE, Schneider SH, Ashe DM, Clark JR, Early R, Etterson JR, Fielder ED, Gill JL, Minter BA, Polasky S, Safford HD, Thompson AR, Vellend M (2009) Multidimensional evaluation of managed relocation. *Proceedings of the National Academy of Sciences of the United States of America* 106: 9721-9724.
- Richardson DM, Hui C, Nuñez MA, Pauchard A (2014) Tree invasions: patterns, processes, challenges and opportunities. *Biological Invasions* 16: 473-481.
- Richardson DM, Le Roux JJ, Wilson JRU (2015) Australian acacias as invasive species: Lessons to be learnt from regions with long planting histories. *Southern Forests* 77: 31-39.
- Richardson DM, Pyšek P, Rejmánek M, Barbour MG, Panetta FD, West CJ (2000) Naturalization and invasion of alien plants: concepts and definitions. *Diversity and Distributions* 6: 93-107.
- Richardson DM, Rejmánek M (2004) Conifers as invasive aliens: a global survey and predictive framework. *Diversity and Distributions* 10: 321-331.
- Richardson DM, van Wilgen BW, Nunez M (2008). Alien conifer invasions in South America – short fuse burning? *Biological Invasions* 10: 573–577.

- Richardson DM, Rejmánek M (2011) Trees and shrubs as invasive alien species - a global review. *Diversity and Distributions* 17: 788-809.
- Riepišas E, Straigitė L (2008) Invasiveness and ecological effects of Red Oak (*Quercus rubra* L.) in Lithuanian forests. *Baltic Forestry* 14(2): 122-130.
- Rodríguez-Loinaz G, Amezaga I, Onaindia M (2013) Use of native species to improve carbon sequestration and contribute towards solving the environmental problems of the timberlands in Biscay, northern Spain. *Journal of Environmental Management* 120: 18-26.
- Romanyà J, Vallejo VR (2004) Productivity of *Pinus radiata* plantations in Spain in response to climate and soil. *Forest Ecology and Management* 195: 177-189.
- Roy et al. 2014 from page 9 is missing here
- Roy HE, Rabitsch W, Scalera R, Stewart A, Gallardo B, Genovesi P, Essl F, Adriaens T, Bacher S, Booy O, Branquart E, Brunel S, Copp GH, Dean H, D'hondt B, Josefsson M, Kenis M, Kettunen M, Linnamagi M, Lucy F, Martinou A, Moore N, Nentwig W, Nieto A, Pergl J, Peyton J, Roques A, Schindler S, Schönrogge K, Solarz W, Stebbing PD, Trichkova T, Vanderhoeven S, van Valkenburg J, Zenetos A (2017) Developing a framework of minimum standards for the risk assessment of alien species. *J Appl Ecol. Ruis BMGS* (2001) No forest convention but ten tree treaties. *Unasylva* 206 – Global Conventions related to forest. An international journal of forestry and forest industries - Vol. 52, 2001/3. FAO - Food and Agriculture Organization of the United Nations, Rome, Italy.
- Sander IL (1990) *Quercus rubra* L. In: Burns R, Honkala B (eds) *Silvics of North America*, Vol. 2. Hardwoods. USDA Agricultural Handbook 654, pp. 1401-1414.
- Sanz-Elorza M, Dana Sánchez ED, Sobrino Vespertinas E (2004) Atlas de plantas alóctonas invasoras en España. Ministerio de Medio Ambiente, Madrid.
- Savill P, Evans J, Auclair D, Falck J (1997) *Plantation Silviculture in Europe*. Oxford University Press. Oxford, reprinted 2005, 297 pp.
- Schiller G, Grunwald C (1987) Resin monoterpenes in rangewide provenance trials of *Pinus halepensis* Mill. in Israel. *Silvae Genetica* 36:109-114.
- Schlyter P, Stjernquist I, Bäckstrand K (2009) Not seeing the forest for the trees? The environmental effectiveness of forest certification in Sweden. *Forest Policy and Economics* 11: 375-382.
- Schmid M, Pautasso M, Holdenrieder O (2014) Ecological consequences of Douglas fir (*Pseudotsuga menziesii*) cultivation in Europe. *European Journal of Forest Research* 133: 13-29.
- Schmithüsen F, Herbst P, Le Master DC (eds.) (2000) *Forging a New Framework for Sustainable Forestry: Recent Developments in European Forest Law*. IUFRO World Series Volume 10. International Union of Forestry Research Organisations, IUFRO Secretariat Vienna; Chair of Forest Policy and Forest Economics, ETH, Zurich, 354 pp.
- Schreck Reis C, Marchante H, Freitas H, Marchante E (2011) Public Perception of Invasive Plant Species: Assessing the impact of workshop activities to promote young students' awareness. *International Journal of Science Education* 35(4): 690-712.
- Searle S, Malins C (2014) A reassessment of global bioenergy potential in 2050. *GCB Bioenergy*. doi: 10.1111/gcbb.12141
- Secretariat of the Convention on Biological Diversity (2001a) Assessment and management of alien species that threaten ecosystems, habitats and species. Abstracts of keynote addresses and posters presented at the sixth meeting of the Subsidiary Body on Scientific, Technical and Technological Advice, held in Montreal, Canada, from 12 to 16 March 2001. Montreal, SCBD, 123p. (CBD Technical Paper no. 1).

- Secretariat of the Convention on Biological Diversity (2001b) Review of the efficiency and efficacy of existing legal instruments applicable to invasive alien species. Montreal, SCBD, 42 pp. (CBD Technical Series no. 2).
- Secretariat of the Convention on Biological Diversity (2002) Review of the status and trends of, and major threats to, the forest biological diversity. Montréal, SCBD. CBD Technical Series no. 7, 164 pp.
- Secretariat of the Convention on Biological Diversity (2009) Sustainable Forest Management, Biodiversity and Livelihoods: A Good Practice Guide. Montreal, 47 + iii pp.
- Sedjo R (2005) Will developing countries be the early adapters of genetically engineered forests? *AbBioForum* 8: 205-212.
- Sein CC, Mitlöhner R (2011) *Acacia* hybrid: ecology and silviculture. CIFOR, Bogor, Indonesia, 13 pp.
- Seo KW, Son Y, Rhoades CC, Noh NJ, Koo JW, Kim J-G (2008) Seedling Growth and Heavy Metal Accumulation of Candidate Woody Species for Revegetating Korean Mine Spoils. *Restoration Ecology* 16: 702-712.
- Shackleton RT, Le Maitre DC, Pasiecznik NM, Richardson DM (2014) *Prosopis*: a global assessment of the biogeography, benefits, impacts and management of one of the world's worst woody invasive plant taxa. *AoB Plants* 6 : plu027 doi:10.1093/aobpla/plu027.
- Shackleton RT, Le Maitre DC, Van Wilgen BW, Richardson DM (2015) The impact of invasive alien *Prosopis* species (mesquite) on native plants in different environments in South Africa. *South African Journal of Botany* 97: 25-31.
- Shine C (2007) Invasive species in an international context: IPPC, CBD, European Strategy on Invasive Alien Species and other legal instruments. *EPPA Bulletin* 37: 103-113.
- Silva JS, Marchante H (2012) Post-fire management of exotic forests. Pages 223-255 in Moreira F, Heras Jdl, Corona P, Arianoutsou M, (eds). *Post-Fire Management and Restoration of Southern European Forests*. Springer, Dordrecht.
- Silvertown J (2009) A new dawn for citizen science. *Trends in Ecology & Evolution* 24: 467-471.
- Simberloff D (2006) Risk assessments, black lists, and white lists for introduced species: are predictions good enough to be useful? *Agricultural and Resource Economics Review* 35: 1-10.
- Simberloff D, Nuñez MA, Ledgard NJ, Pauchard A, Richardson DM, Sarasola M, Van Wilgen BW, Zalba SM, Zenni RD, Bustamante R, Peña E, Ziller SR (2010) Spread and impact of introduced conifers in South America: Lessons from other southern hemisphere regions. *Austral Ecology* 35: 489-504.
- Singh AN, Zeng DH, Chen FS (2006) Effect of young woody plantations on carbon and nutrient accretion rates in a redeveloping soil on coalmine spoil in a dry tropical environment, India. *Land Degradation & Development* 17: 13-21.
- Siple MC, Donahue MJ (2013) Invasive mangrove removal and recovery: Food web effects across a chronosequence. *Journal of Experimental Marine Biology and Ecology* 448: 128-135.
- Sitzia T, Campagnaro T, Dainese M, Cierjacks A (2012) Plant species diversity in alien black locust stands: a paired comparison with native stands across a North-Mediterranean range expansion. *Forest Ecology and Management* 285: 85-91.
- Sitzia T, Trentanovi G., Marini L, Cattaneo D, Semenzato P. (2013) Assessment of hedge stand types as determinants of woody species richness in rural field margins. *iForest-Biogeosciences and Forestry*, 6(4): 201-208.
- Sitzia T (2014) A call to silviculturists for a new field of science: The forestry of invasive alien species. *The forestry chronicle* 90(4): 486-488.

- Slade R, Bauen A, Gross R (2014) Global bioenergy resources. *Nature Climate Change*. doi:10.1038/nclimate2097.
- Sladonja B, Sušek M, Guillermic J (2015) Review on invasive tree of Heaven (*Ailanthus altissima* (Mill.) Swingle) conflicting values: assessment of its ecosystem services and potential biological threat. *Environmental Management* 56(4): 1009-1034.
- Slavov GT, Leonardi S, Burczyk J, Adams WT, Strauss SH, DiFazio SP (2009) Extensive pollen flow in two ecologically contrasting populations of *Populus trichocarpa*. *Molecular Ecology* 18: 357-373.
- SLU (2010) Skogsdata 2010, Sveriges officiella statistik, Umeå, 115 pp. (In Swedish with English summaries) [http://pub.epsilon.slu.se/5421/1/Skogsdata2010_webb.pdf].
- Smulders M, Beringen R, Volosyanchuk R, Vanden Broeck A, Schoot J, Arens PFP, Vosman B (2008) Natural hybridisation between *Populus nigra* L. and *P. x canadensis* Moench. Hybrid offspring competes for niches along the Rhine river in the Netherlands. *Tree Genetics & Genomes* 4: 663-675.
- Somerville C, Youngs H, Taylor C, Davis SC, Long SP (2010) Feedstocks for Lignocellulosic Biofuels. *Science* 329: 790-792. doi: 10.1126/science.1189268
- Sserwanga A, Harris JC, Kigozi R, Menon M, Bukirwa H, Gasasira A, Kakeeto S, Kizito F, Quinto E, Rubahika D, Nasr S, Filler S, Kanya MR, Dorsey G (2011) Improved malaria case management through the implementation of a health facility-based sentinel site surveillance system in Uganda. *PLoS One* 6:e16316
- Stanturf JA, Palik BJ, Dumroese RK (2014) Contemporary forest restoration: A review emphasizing function. *Forest Ecology and Management* 331: 292-323.
- Starfinger U (1997) Introduction and naturalization of *Prunus serotina* in Central Europe. In: Brock J, Wade M, Pysek P, Green D (eds.), *Plant invasions: studies from North America and Europe*. Leiden: Backhuys Publishers; The Netherlands, pp. 161-171.
- Starfinger U, Kowarik I (2003) *Prunus serotina* Ehrh. (Rosaceae), Späte Traubenkirsch. NeoXora fact sheet (<http://www.floraweb.de/neoflora/handbuch/prunusserotina.pdf>).
- Starfinger U, Kowarik I, Rode M, & Schepker H (2003) From desirable ornamental plant to pest to accepted addition to the flora? – the perception of an alien tree species through the centuries. *Biological Invasions* 5: 323-335.
- Starfinger U (2010) NOBANIS – Invasive Alien Species Fact Sheet – *Prunus serotina* – From: Online Database of the European Network on Invasive Alien Species – NOBANIS www.nobanis.org [Accessed 30 August 2015].
- Strassburg BBN, Kelly A, Balmford A, Davies RG, Gibbs HK, Lovett A, Miles L, Orme CDL, Price J, Turner RK, Rodrigues ASL (2010) Global congruence of carbon storage and biodiversity in terrestrial ecosystems. *Conservation Letters*, 3: 98-105.
- Strauss SH, Brunner AM, Busov VB, Ma CP, Meilan R (2004). Ten lessons from 15 years of transgenic *Populus* research. *Forestry* 77: 455-465.
- Strauss SH, Tan H, Boerjan W, Sedjo R (2009) Strangled at birth? Forest biotech and the Convention on Biological Diversity. *Nature Biotechnology* 27: 519-527.
- Stromberg JC, Lite SJ, Marler R, Paradzick C, Shafroth PB, Shorrock D, White JM, White MS (2007) Altered stream-flow regimes and invasive plant species: the *Tamarix* case. *Global Ecology and Biogeography*, 16: 381-393.
- Stupak I, Lattimore B, Titus BD, Tattersall Smith C (2011) Criteria and indicators for sustainable forest fuel production and harvesting: A review of current standards for sustainable forest management. *Biomass and Bioenergy* 35: 3287-3308.

- Sturgess P, Atkinson D (1993) The clear-felling of sand-dune plantations: soil and vegetational processes in habitat restoration. *Biological Conservation* 66: 171-183.
- Suzuki N, Olson DH (2008) Options for biodiversity conservation in managed forest landscapes of multiple ownerships in Oregon and Washington, USA. *Biodiversity and Conservation* 17: 1017-1039.
- Szítár K, Ónodi G, Somay L, Pándi I, Kucs P, Kröel-Dulay G (2014) Recovery of inland sand dune grasslands following the removal of alien pine plantation. *Biological Conservation* 171: 52-60.
- Taylor V, Kecse-Nagy K, Osborn T (2012) Trade in *Dalbergia nigra* and the European Union. Report prepared for the European Commission, 24 pp. [http://ec.europa.eu/environment/cites/pdf/Dalbergia%20Report_FIN%2020%2012%202012.pdf].
- Thompson I, Mackey B, McNulty S, Mosseler A (2009) Forest Resilience, Biodiversity, and Climate Change. A synthesis of the biodiversity/resilience/stability relationship in forest ecosystems. Secretariat of the Convention on Biological Diversity, Montreal. Technical Series no. 43, 67 pp.
- Tokarska-Guzik B, Dajdok Z, Zajac M, Zajac A, Urbisz A, Danielewicz W, Hołdyński C (2012) Rośliny obcego pochodzenia w Polsce ze szczególnym uwzględnieniem gatunków inwazyjnych, Warsaw, 107 pp. (In Polish).
- Todorović S, Božić D, Simonović A, Filipović B, Dragičević M, Giba Z, Grubišić D (2010) Interaction of fire-related cues in seed germination of the potentially invasive species *Paulownia tomentosa* Steud. *Plant Species Biology* 25: 193-202.
- Trinka P (1998) Short-rotation forestry: discussion of 10 Austrian principles from the viewpoint of preservation of environment and nature. *Biomass and Bioenergy* 15(1): 109-114.
- UNCED (1992a) Non-Legally Binding Authoritative Statement of Principles for a Global Consensus on the Management, Conservation and Sustainable Development of all Types of Forests, UN A/CONF.151/26 (Vol. III).
- UNCED (1992b) Agenda 21 Chapter 11. In: *Combating Deforestation*, Rio de Janeiro.
- UNCED (1992c) Forest principles. In: *Report of the United Nations Conference on Environment and Development*, Rio de Janeiro, 3–14 June.
- USDA (2012) Non-native invasive species best management practice. Guidance for the US Forest Service Eastern Region. United States Department of Agriculture, Forest Service, August 2012, 282 pp.
- Van den Meersschaut D, Lust N (1997) Comparison of mechanical, biological and chemical methods for controlling black cherry (*Prunus serotina*) in Flanders (Belgium). *Silva Gandavensis* 62: 90-109.
- Van Wilgen BW, Richardson DM (2012) Three centuries of managing introduced conifers in South Africa: benefits, impacts, changing perceptions and conflict resolution. *Journal of Environmental Management* 106: 56-68.
- Van Wilgen BW, Richardson DM (2014) Challenges and trade-offs in the management of invasive alien trees. *Biological Invasions* 16: 721-734.
- Van Wyk DB (1987) Some effects of afforestation on streamflow in the Western Cape Province, South Africa. *Water SA* 13: 31-36.
- Vanden Broeck A, Villar M, van Bockstaele E, van Slycken J (2005) Natural hybridization between cultivated poplars and their wild relatives: evidence and consequences for native poplar populations. *Annals of Forest Science* 62: 601-613.
- Vanhellemont M, Verheyen K, Staelens J, Hermy M (2010) Factors affecting radial growth of the invasive *Prunus serotina* in pine plantations in Flanders. *European Journal of Forest Research* 129(3): 367-375.

- Venette RC, Kriticos DJ, Magarey RD, Koch FH, Baker RHA, Worner SP, Gómez Raboteaux NN, McKenney DW, Dobesberger EJ, Yemshanov D, De Barro PJ, Hutchison WD, Fowler G, Kalaris TM, Pedlar J (2010) Pest risk maps for invasive alien species: a roadmap for improvement. *BioScience* 60(5): 349-362.
- Verheyen K, Vanhellemont M, Auge H, Baeten L, Baraloto C, Barsoum N, Bilodeau-Gauthier S, Bruelheide H, Castagneyrol B, Godbold D, Haase J, Hector A, Jactel H, Koricheva J, Loreau M, Mereu S, Messier C, Muys B, Nolet P, Paquette A, Parker J, Perring M, Ponet Q (2016) Contributions of a global network of tree diversity experiments to sustainable forest plantations. *Ambio* 45: 29.
- Vettrano AM, Roques A, Yart A, Fan J-t, Sun H-h, Vannini A (2015) Sentinel trees as a tool to forecast invasions of alien plant pathogens. *PloS One*, doi: 10.1371/journal.pone.0120571.
- Visser V, Langdon B, Pauchard A, Richardson DM (2014) Unlocking the potential of Google Earth as a tool in invasion science. *Biological Invasions* 16: 513-534.
- Wakie TT, Evangelista PH, Jarnevich CS, Laituri M (2014) Mapping Current and Potential Distribution of Non-Native *Prosopis juliflora* in the Afar Region of Ethiopia. *PLoS One* 9(11): e112854. doi:10.1371/journal.pone.0112854.
- Wasowicz P, Przedpelska-Wasowicz E, Kristinsson H (2013) Alien vascular plants in Iceland: diversity, spatial patterns, temporal trends, and the impact of climate change. *Flora* 208: 648-673.
- Weih M (ed) (2008) Short rotation forestry (SRF) on agricultural land and its possibilities for sustainable energy supply. *TemaNord* 2008:543, Nordic Council of Ministers, Copenhagen [www.norden.org/publications].
- Werner C, Zumkier U, Beyschlag, W, Máguas C (2010) High competitiveness of a resource demanding invasive acacia under low resource supply. *Plant Ecology* 206: 83-96.
- Westbrooks R (2003) A National Early Detection and Rapid Response System for Invasive Plants in the United States: Conceptual Design. Federal Interagency Committee for the Management of Noxious and Exotic Weeds (FICMNEW).
- Widrlechner MP, Thompson JR, Iles JKD, Dixon PM (2004) Models for predicting the risk of naturalization of nonnative woody plants in Iowa. *Journal of Environmental Horticulture* 22: 23-31.
- Williams MRW, Winn M (1977) A plantation of monkey puzzle at Monreith, Wigtownshire. *Scottish Forestry* 31(2): 71-73.
- Wilson JRU, Caplat P, Dickie IA, Hui C, Maxwell BD, Nuñez MA, Pauchard A, Rejmánek M, Richardson DM, Robertson MP, Spear D, Webber BL, van Wilgen BW, Zenni RD (2014) A standardized set of metrics to assess and monitor tree invasions. *Biological Invasions* 16: 535-551.
- Wilson JRU, Dormontt EE, Prentis PJ, Lowe AJ, Richardson DM (2009) Something in the way you move: dispersal pathways affect invasion success. *Trends in Ecology & Evolution* 24: 136-144.
- Wingfield MJ, Brockerhoff EG, Wingfield BD, Slippers B (2015) Planted forest health: The need for a global strategy. *Science* 349 (6250): 832-836.
- Wise RM, van Wilgen BW, Le Maitre DC (2012) Costs, benefits and management options for an invasive alien tree species: The case of mesquite in the Northern Cape, South Africa. *Journal of Arid Environments* 84: 80-90.
- Wishnie MH, Dent DH, Mariscal E, Deago J, Cedeno N, Ibarra D, Condit R, Ashton PMS (2007) Initial performance and reforestation potential of 24 tropical tree species planted across a precipitation gradient in the Republic of Panama. *Forest Ecology and Management* 243: 39-49.

- Witt ABR (2010) Biofuels and invasive species from an African perspective - a review. *GCB Bioenergy* 2: 321-329.
- Wolfslehner B, Vacik H, Lexer MJ (2005) Application of the analytic network process in multi-criteria analysis of sustainable forest management. *Forest Ecology and Management* 207: 157–170.
- Woziwoda B, Kopeć D, Witkowski J (2014) The negative impact of intentionally introduced *Quercus rubra* L. on a forest community. *Acta Societatis Botanicorum Poloniae* 83(1): 39-49.
- Wu P-F, Ma X-Q, Tigabu M, Huang Y, Zhou L-L, Cai L, Hou X-L, Oden PC (2014) Comparative growth, dry matter accumulation and photosynthetic rate of seven species of *Eucalypt* in response to phosphorus supply. *Journal of Forestry Research* 25(2): 377-383.
- Yelenik SG, Stock WD, Richardson DM (2004) Ecosystem level impacts of invasive *Acacia saligna* in the South African fynbos. *Restoration Ecology* 12: 44-51.
- Yelenik SG, Stock WD, Richardson DM (2007) Functional group identity does not predict invader impacts: differential effects of nitrogen-fixing exotic plants on ecosystem function. *Biological Invasions* 9: 117-125.
- Zapponi L, Minari E, Longo L, Toni I, Mason F, Campanaro A (2014) The Habitat-Trees experiment: using exotic tree species as new microhabitats for the native fauna. *iForest-Biogeosciences and Forestry*, 8: 464-470.

6. ANNEXES

6.1 Definitions – Glossary

The terminology used in legislation and in the scientific and technical literature when discussing alien trees and invasive tree species can be complex and confusing as many of the terms have been used in different ways by different authors. Unless referenced, definitions follow FAO, CBD, Bern Convention, Regulation EU no. 1143/2014, the Code of Conduct on Horticulture and Invasive Alien Plants (Heywood & Brunel 2009, 2011) and the European Code of Conduct for Botanic Gardens on Invasive Alien Species (Heywood & Sharrock 2013).

Alien tree species

A tree species, subspecies or lower taxon, introduced outside its natural past or present distribution; includes any part, gametes, seeds, or propagules of such species that might survive and subsequently reproduce (Decision VI/23 of the Conference of the Parties to the CBD, Annex, footnote to the Introduction).

In the context of the present Code of Conduct the terms alien, non-native, exotic and introduced tree are considered as equivalent. In accordance with the CBD definition, the term alien tree has exclusively a biogeographical meaning, i.e. it refers to a species, subspecies or lower taxon, introduced outside its natural past or present distribution; it includes any part, seeds, or propagules of such species that might survive and subsequently reproduce. As such, the term alien tree does not include any negative evaluation of the tree species. Only a small percentage of all the alien tree species are, or may become after some time, invasive alien tree species (COP 6 Decision VI/23 “Alien species that threaten ecosystems, habitats or species”). Importantly, an alien tree species is “introduced outside its natural past or present distribution” deliberately or accidentally by man. The definition takes into consideration the Recommendation No. 142 (2009) of the Standing Committee (Convention on the Conservation of European Wildlife and Natural Habitats), adopted on 26 November 2009, interpreting the CBD definition of invasive alien species to take into account climate change, “recommends Contracting Parties to the Convention and invites Observer States to: 1. interpret the term “alien species” for the purpose of the implementation of the European Strategy on Invasive Alien Species as **not including native species naturally extending their range in response to climate change**” (Cf. Section 4.6.2 in this Code). As a result, also past mass migratory events in forest tree populations, postglacial recolonization routes and similar events are not considered herewith in the definition of alien tree species. We focus on alien trees deliberately or accidentally introduced by man outside its natural past or present distribution, where “past” refers to the definition of “neophytes” (i.e. introduced after the 1,500) as used in the CBD context and defined by Pyšek et al. (2004). In addition, with specific concern to this Code, an alien tree species is alien to all the 47 Member States of the Council of Europe. According to this definition the term 'alien tree species' does not include foreign provenances of tree species that are native in at least one of the 47 Member States of the Council of Europe.

Cf. also the Regulation (EU) No. 1143/2014 of the European Parliament and of the Council, of 22 October 2014, on the prevention and management of the introduction and spread of invasive alien species (Article 3 – Definitions: 'alien species' means any live specimen of a species, subspecies or lower taxon of animals, plants, fungi or micro-organisms introduced outside its natural range; it includes any part, gametes, seeds, eggs or propagules of such species, as well as any hybrids, varieties or breeds that might survive and subsequently reproduce).

Afforestation

Afforestation is the act of establishing forests through planting and/or deliberate seeding on land that, until then, was not classified as forest (FAO 2010a, 2015a, 2015b). Reforestation on the other hand, takes place in areas that already are classified as forest and does not imply any change of land use from a non-forest use to forest.

Ancient forest

An ancient forest is a forest that has existed continuously since at least a specified date (threshold date), selected on the availability of historical land-use information and differing between studies and countries (Hermy et al. 1999; Verheyen & Hermy 2007).

Bioenergy

Bioenergy¹⁰⁹ is the conversion of biomass resources into useful energy carriers including heat, electricity and transport fuels. Biomass is derived from different types of organic matter: energy plants (oilseeds, plants containing sugar) and forestry, agricultural or urban waste including wood and household waste. Biomass can be used for heating, for producing electricity and for transport biofuels. Biomass can be solid (plants, wood, straw and other plants), gaseous (from organic waste, landfill waste) or liquid (derived from crops such as wheat, rapeseed, soy, or from lignocellulosic material).

Black List

A Black List identifies those alien species whose introduction is prohibited due to their potential adverse effects on the environment or human, animal or plant health. Such lists can be a significant component of an invasive alien species prevention regime since they clearly state which species are banned from import. Black lists are the most common type of listing mechanism and are found in a range of countries. Such lists are most useful to prevent intentional introductions at the pre-border stage, as a potential exporter can check the relevant lists to see if the species in question is allowed or banned from imports, or, for unlisted species, request permission to import. This provides increased transparency and predictability for exporters before any products are gathered, packaged and shipped. Lists can also be used at the border by inspection and quarantine agents for purposes of searching baggage, package and cargo.

The success of such a listing system is inherently related to its adaptability and flexibility, particularly with regard to processing new submissions and proposals for movement from one list to another. The three types of lists are referred to as black, white and grey lists, and are sometimes used individually and sometimes in combination. More in general, black lists are nowadays not only restricted to the pre-import stage.

CBD - Convention on Biological Diversity

Signed by 150 government leaders at the 1992 Rio Earth Summit, the Convention on Biological Diversity is dedicated to promoting sustainable development. Conceived as a practical tool for translating the principles of Agenda 21 into reality, the Convention recognizes that biological diversity is about more than plants, animals and microorganisms and their ecosystems – it is about people and our need for food security, medicines, fresh air and water, shelter, and a clean and healthy environment in which to live (<http://www.cbd.int/convention/default.shtml>).

Eradication

The extirpation of an entire population of an alien species within a designated management unit. When a species can be declared eradicated (that is, how long a period of time after the management intervention) depends on the species and the situation and must take into account factors such as seed-bank longevity (for plants). Eradication success should be stated in terms of confidence limits (e.g. 1-5 % confidence) that the species is not present. Eradication is possible in many cases, but there are no clearly documented cases of the eradication of an alien tree species (van Wilgen & Richardson 2014).

¹⁰⁹ http://ec.europa.eu/energy/renewables/bioenergy/bioenergy_en.htm

FRA – Forest Resources Assessment

FAO has been monitoring the world's forests at 5 to 10-year intervals since 1946. The Global Forest Resources Assessments (FRA) are now produced every five years in an attempt to provide a consistent approach to describing the world's forests and how they are changing, e.g. FRA 2010, 2015. The scope and content of the global assessments have evolved over time to respond to changing information needs (FAO 2015a, 2015b).

Impact

The description or quantification of how an invasive alien tree affects the physical, chemical and biological environment. Many invasive alien tree species have substantial effects on the ecosystems into which they have been introduced, including significant changes in native species extinction probabilities, genetic composition, behaviour patterns, richness and abundance, as well as altering phylogenetic and taxonomic diversity, trophic networks, ecosystem productivity, nutrient cycling, geomorphology, hydrology, habitat structure and various components of disturbance regimes (Hawkins et al. 2015).

Invasive alien tree species

In accordance with the CBD definition, and for the purposes of this Code, an **invasive alien tree** is herewith defined as an alien tree species whose introduction and/or spread threaten or adversely impact upon biodiversity and related ecosystem services.

Not all alien trees will become naturalised. Similarly, **invasive alien trees** are a subset of the naturalised alien trees, as many naturalised alien trees do not go on to become invasive, i.e. they do not threaten biological diversity and do not cause economic or environmental harm.

Cf. also the Regulation (EU) No. 1143/2014 of the European Parliament and of the Council, of 22 October 2014, on the prevention and management of the introduction and spread of invasive alien species (Article 3 – Definitions: 'invasive alien species' means an alien species whose introduction or spread has been found to threaten or adversely impact upon biodiversity and related ecosystem services).

For example, according to FAO (2012), **woody invasive species** are woody plant species that are non-native to a particular ecosystem and whose introduction and spread cause, or are likely to cause, socio-cultural, economic or environmental harm or harm to human health.

The FAO 2015 Global Forest Resources Assessment (FRA), defines an **introduced species** any taxon (from genus to sub-species) occurring outside its natural range (past or present) and out of its dispersal potential area, i.e. outside the range it occupies naturally or could occupy without direct or indirect introduction or care by humans, FAO (2012).

In FRA 2015, the term introduced is considered equivalent to non-native. In this Code the terms introduced, non-native, and exotic are considered equivalent to alien. In addition, in FAO FRA 2015, “**naturalised tree**” species are introduced tree species that spread and multiply by natural regeneration and are well established and acclimatized for several years. They do not need human help to reproduce/maintain populations over time (FAO 2012).

Tree invasiveness

The features of invasive alien tree species, such as their life-history traits and modes of reproduction, that define their capacity to invade, i.e. to overcome various barriers to invasion and produce negative impacts. The level of invasiveness of a species can change over time due to, for example, changes in genetic diversity through hybridization, introgression, or the continued arrival of new propagules of the same species that is already established in a region, but from new and different (meta)populations, such that genetic diversity may increase.

Management of invaded habitat as a “novel ecosystem”

Ensuring the continued and sustainable delivery of key functions and services, in some cases accepting that invasive alien species fulfil useful purposes, especially where conditions are modified to the extent that the return of native species is unrealistic (van Wilgen & Richardson 2014). Where habitats have been substantially modified through multiple human factors, removing invasive alien trees and restoring native dominated communities and ecosystem functions is sometimes either impossible or undesirable. For instance, in riparian ecosystems in many parts of the world that are heavily invaded by alien trees, physical conditions have been modified to such an extent that native elements can no longer establish or survive, even when the invasive trees are removed. In such cases, manipulating of the density and abundance of key alien species to achieve desired ecosystem functions and services is an appropriate, pragmatic management goal (Richardson et al. 2007).

Pest

According to the International Plant Protection Convention (IPPC) a pest is “any species, strain or biotype of plant, animal or pathogenic agent injurious to plants or plant products”, while a quarantine pest is “a pest of potential economic importance to the area endangered thereby and not yet present there, or present but not widely distributed and being officially controlled”. As a consequence, considering that potential economic importance can account for environmental concern (according to the supplement the International Standard on Phytosanitary Measures No. 5 Glossary of phytosanitary terms), the IPPC definition of a quarantine pest covers much of what is considered an invasive alien species under the CBD. Differences arise from the fact that a quarantine pest does not necessarily have to be alien, threaten biodiversity, may only affect agriculture, and that an invasive alien plant may not be considered a quarantine pest if it is widely distributed (Heywood & Sharrock 2013). *Cf* also, Commission Implementing Directive (EU) 2017/1279 of 14 July 2017 amending Annexes I to V to Council Directive 2000/29/EC on protective measures against the introduction into the Community of organisms harmful to plants or plant products and against their spread within the Community.

Polluter Pays Principle (PPP)

The Polluter-Pays Principle (PPP) was adopted by The Organisation for Economic Co-operation and Development in 1972 as an economic principle for allocating the costs of pollution control. Under the 1972 and 1974 OECD Recommendations, the Polluter-Pays Principle means that the polluter should bear the "costs of pollution prevention and control measures", the latter being "measures decided by public authorities to ensure that the environment is in an acceptable state". The principle is laid down in the Rio Declaration (CBD) and in Directive 2004/35/EC of the European Parliament and of the Council of 21 April 2004 on environmental liability.

Reforestation

Reforestation refers to the re-establishment of forest through planting and/or deliberate seeding on land classified as forest, for instance after a fire, storm or following clearfelling (FAO 2010a).

Residence time

The time since the introduction of an alien species to a region; since the introduction date is usually derived from post-hoc records and is likely inaccurate, the term minimum residence time has been suggested. The extent of invasion of alien species generally increases with increasing residence time as species have more time to fill their potential ranges.

Risk assessment (RA)

The estimation of the quantitative or qualitative value of risk (the likelihood of an event occurring within a specified time frame and the consequences if it occurs). In the context of invasion ecology, RA is undertaken to evaluate the likelihood of the entry, establishment and spread of an alien species (intentionally or accidentally) in a given region, negotiating given barriers in the naturalization-invasion

continuum, and the extent and severity of ecological, social and economic impacts. Risk assessment is defined by article 5 of the WTO Agreement on the Application of Sanitary and Phytosanitary Measures (SPS Agreement). Risk assessment is one of the three components of Risk Analysis (risk assessment, risk management and risk communication).

6.2 Alien tree species listed in European countries

Species	DK	BL	IT	IR	Malta	NW	PT	S W	EPPO
<i>Abies alba</i>						HI			Invasive
<i>Abies balsamea</i>						LO			
<i>Abies concolor</i>						PH			
<i>Abies grandis</i>						PH			
<i>Abies koreana</i>						LO			
<i>Abies lasiocarpa</i>						LO			
<i>Abies mariesii</i>						NK			
<i>Abies procera</i>						LO			
<i>Abies sibirica</i>						PH			
<i>Acacia cyanophylla</i>							Annex I		
<i>Acacia cyclops</i>					MPI				
<i>Acacia dealbata</i>				Potential			Annex I	Watch- List	Invasive
<i>Acacia farnesiana</i>							Annex I-III		
<i>Acacia karroo</i> (= <i>Vachellia karroo</i>)					MPI		Annex I		
<i>Acacia longifolia</i>							Annex I		
<i>Acacia mearnsii</i>							Annex I		
<i>Acacia melanoxylon</i>				Potential			Annex I		
<i>Acacia pycnantha</i>							Annex I		
<i>Acacia retinodes</i>							Annex I		
<i>Acacia saligna</i>					MPI				
<i>Acer campestre</i>						LO			
<i>Acer negundo</i>		WL B2	Black- List			LO			
<i>Acer platanoides</i>				Uncertain					
<i>Acer pseudoplatanus</i>				Amber- List		SE			
<i>Aesculus hippocastanum</i>						PH			
<i>Ailanthus altissima</i>		BL A2	Black- List	Uncertain	MPI		Annex I	Black-List	Invasive
<i>Casuarina equisetifolia</i>					MPI				
<i>Chamaecyparis lawsoniana</i>						LO			
<i>Eucalyptus camaldulensis</i>					MPI				
<i>Larix decidua</i>						SE			
<i>Paulownia tomentosa</i>			Black- List	Potentia l				Watch- List	
<i>Picea sitchensis</i>				Uncerta in		SE			
<i>Pinus contorta ssp. contorta</i> var. <i>contorta</i>	Black- List			Uncerta in		PH			
<i>Pinus contorta ssp. contorta</i> var. <i>latifolia</i>	Black- List								
<i>Pinus contorta ssp. murrayana</i>	Black- List								
<i>Pinus mugo ssp. mugo</i>	Black- List					SE			
<i>Pinus mugo ssp. mugo</i> x <i>rotundata</i>	Black- List								
<i>Pinus nigra</i>			Black- List			LO			

<i>Populus alba</i>					LO		
<i>Prunus laurocerasus</i>		WL	Black-				Black-
		B1	List				List
<i>Prunus serotina</i>	Black-	BL	Black-		HI		Black-
	List	A3	List				List
<i>Pseudotsuga menziesii</i>				Uncerta	LO		
				in			
<i>Quercus cerris</i>				Uncerta	LO		
				in			
<i>Quercus rubra</i>		WL	Black-	Uncerta	LO		
		B3	List	in			
<i>Robinia pseudoacacia</i>	Obs-	WL	Black-	Uncerta	HI	Annex	*
	List	B3	List	in		I	Black-
							List
<i>Rhus typhina</i>		WL			NK		Black-
		B1					List
<i>Salix viminalis</i>				Uncerta	PH		
				in			
<i>Thuja plicata</i>					LO		

Table 1 - The alien trees most frequently listed (within different categories) in different European countries (DK = Denmark, BL = Belgium, IT = Italy, IR = Ireland, NW = Norway, PT = Portugal, SW = Switzerland, EPPO = EPPO Region). The table includes both tree species alien “to” and alien “in” Europe or in the EPPO region. The term “alien in” is used for those tree species that are native in some European countries but are considered alien in other European countries (e.g. *Populus alba*). Plants names are reported exactly as they are found in the original source, regardless of synonyms or invalid names (e.g. *Acacia cyanophylla* Lindl. is a synonym for *Acacia saligna* (Labill.) H.L.Wendl.; *Pinus contorta* ssp. *murrayana* should be indicated as *Pinus contorta* var. *murrayana*). * see comment below in the paragraph on Switzerland.

In **Denmark**, non-native species are not explicitly dealt with under the Forestry Act, but through some of the statutory orders affiliated with this law various lists of accepted trees/shrubs are maintained by the Danish Nature Agency (Madsen et al. 2014).

The **Belgian** Forum on Invasive Species (<http://ias.biodiversity.be/>) provides information on Alert, Black and Watch lists of invasive species in Belgium on its web site (Branquart 2014).

In **Italy** two regional Black Lists are in force, i.e. in Lombardy (LR no. 10, 31 March 2008; DGR VIII/007736, 24 July 2008) and Piedmont (Determinazione Regionale DB0701 no. 448, 25 May 2012; DGR 46-5100, 18 December 2012) (Brundu 2008).

Invasive Species **Ireland**, a joint venture between the Northern Ireland Environment Agency and the National Parks and Wildlife Service (<http://invasivespeciesireland.com/>), produces lists of invasive and non-native species in Ireland and Northern Ireland using the Non-native species APplication based Risk Analysis (NAPRA).

The **Malta**¹¹⁰ Environment and Planning Authority (MEPA, <http://www.mepa.org.mt/>) has commissioned two studies to list alien plant and animal species found in the Maltese Islands and to

¹¹⁰ Under Part III of the “Trees and Woodlands Protection Regulations, 2011” (Legal Notice 200 of 2011) the species included in Schedule III are deemed to be species causing damage to biological diversity of trees or woodlands in Malta, or to the natural environment in general (Regulation 8, paragraph 1). The species in question are *Acacia cyclops*, *Acacia saligna* [= *Acacia cyanophylla*], *Vachellia karroo* [= *Acacia karroo*], *Ailanthus altissima*, *Eucalyptus camaldulensis*, *Eucalyptus gomphocephala*, *Leucaena leucocephala* [= *Acacia leucocephala*; *Albizia lebbek*], *Pittosporum tobira*, *Ricinus communis* and *Schinus terebinthifolius*. Regulation 8, paragraph 2, prohibits the propagation, sowing, import, export, transportation, selling or exchanging any of these species. Regulation 9 on improvement measures also applies to the species listed in Schedule III of Legal Notice 200 of 2011. Apart from these species, Regulation 10 enables the Competent Authority to stop the transport and importation of trees, which may endanger the biological diversity of trees or woodlands in Malta, or other reasons as stated in the provision. In this respect, it should be noted that these species listed in the Regulations have all been proven to have an adverse

identify the invasive types which require further action such as eradicating or controlling their spread in protected areas.

The **Norwegian** Biodiversity Information Centre is responsible for assessing the ecological impacts associated with species that are non-native to Norway (alien species) and to provide an overview of alien species found in Norway (Gederaas et al. 2012).

In 1999, the **Portuguese** legislation addressed the problem of invasive alien species with the Decreto-Lei no. 565/99, of the 21st December 1999, which regulates the introduction of non-native species. This law lists the introduced alien species in Portugal, indicating which are considered invasive and prohibiting the introduction of new species (with some exceptions). Furthermore, the legislation prohibits the possession, cultivation, growing and the trade of species that are considered invasive or of ecological risk (<http://invasoras.pt/en/in-portugal/>). With concern to *Robinia pseudoacacia*, the Decree-Law No. 205/2003, of 12 September, transposes into national law the Council Directive 1999/105/EC of December 22, on the marketing of forest reproductive material, and partially repeals the provisions of Article 8 paragraph 2 of Decree-Law No. 565/99, of 21 December, in that it establishes the prohibition of transfer, purchase, sale, offering for sale and transport of live specimens, as well as the production for trade of the same species. However, *Robinia pseudoacacia* continues to be banned for use in Portugal.

impact on Maltese biodiversity. The Regulations are available at: <http://www.justiceservices.gov.mt/DownloadDocument.aspx?app=lom&itemid=11493&l=1>

Afforestation (and deforestation) is also included in Schedule IA, that is “Projects which require an Environmental Impact Statement or an Environmental Planning Statement”, in the “Environmental Impact Assessment Regulations, 2007, as amended” (Legal Notice 114 of 2007, as amended by Legal Notices 425 of 2007, 438 of 2011 and 211 of 2015). These are available at: <http://www.justiceservices.gov.mt/DownloadDocument.aspx?app=lom&itemid=11556&l=1>

In 2002, the then Planning Authority (now MEPA) published the “Guidelines on Trees, Shrubs and Plants for Planting and Landscaping in the Maltese Islands”, available at: <https://www.mepa.org.mt/LpDocumentDetails?syskey=244>. The 2002 guidelines aim to: (1) promote environmentally-sound planting and soft-landscaping by guiding interested agencies (e.g. Government Departments, Local Councils, voluntary organisations) and the general public; (2) encourage incentives for environmentally-compatible improvements in planting and landscaping projects, and to deter unsustainable, or environmentally-damaging practice; (3) further promote the demand for the propagation of suitable indigenous vegetation, and encourage Governmental and private nurseries to satisfy such demand; and (4) enable clients/developers, as well as their architects and consultants, to produce appropriate landscaping layouts and drawings for specific development projects. Appendix V to the 2002 guidelines lists those species that are unacceptable in rural areas. This list includes tree species that are invasive in the Maltese Islands. More recently, MEPA also published in 2009, the illustrated booklet (not available in digital format) entitled “Common Species used for Landscaping in the Maltese Islands”. This booklet covers soft landscaping (interventions based on planting) and classifies the plant species that are illustrated into the following categories: (a) plants that are acceptable to use in landscaping: large trees; smaller trees and larger shrubs; and smaller shrubs and creepers; (b) trees for particular locations; (c) alien species.

Within the context of invasive alien plants, MEPA adopted on 7 March 2013 the publication entitled “Guidelines on managing non-native plant invaders and restoring native plant communities in terrestrial settings in the Maltese Islands” available at: <http://www.mepa.org.mt/guidelines-alienplants>. The purpose of these guidelines is to: (1) assist the planning and implementation of management programmes, aimed at counteracting the spread of existing plant invaders in important natural and semi-natural areas as well as rural areas, where the removal of non-native plants is desired; and (2) assist the design and implementation of native plant conservation translocations (such as plant reintroductions or reinforcements), aimed at reinstating native plant communities to a favourable conservation status or reinstate an ecological function. The document also serves as guidance to be followed when drawing up method statements on the removal of invasive plants and when implementing conditions that accompany development permits. The invasive woody species (among other plants) addressed in the Guidelines are: *Acacia cyclops*, *Vachellia karroo*, *Acacia saligna*, *Ailanthus altissima*, *Casuarina equisetifolia*, *Eucalyptus* spp., *Leucaena leucocephala*, *Nicotiana glauca*, *Ricinus communis* and *Schinus terebinthifolius*.

The **Swiss** "Ordinance on the Handling of Organisms in the Environment (Release Ordinance, RO)" is the legal basis for the handling of organisms in the environment (CC 814.911 - <https://www.admin.ch/opc/en/classified-compilation/20062651/index.html>). Invasive alien organisms, in accordance with Annex 2 of this ordinance, may not be handled directly in the environment, other than in the case of measures to control them. Other important references are the Law for the protection of the Environment (LPAmb, RS 814.01, 7 October 1983) and the Federal Law on the protection of Nature and Landscape (LPN, RS 451, 1st July 1966). The Swiss Forest Act on Forest (ForA, RS 821.0) regulates the forests and forestry activities. *The federal ordinance on forestry reproduction material (RS 921.552.1) lists and regulates forest tree species, including a number of alien species allowed under certain circumstances in forestry activities. The federal Plant protection ordinance, OPV, RS 916.20 regulates pests.

The panel of experts on invasive alien plants (European and Mediterranean Plant Protection Organisation, **EPPO**) has established lists of Invasive Alien Plants (EPPO A1/A2 list, List of invasive alien plants, Observation List and Alert list)¹¹¹ on the basis of transparent criteria and using the EPPO Prioritization Process on Invasive Alien Plants. EPPO recommends countries endangered by these species to consider measures to prevent their introduction and spread or to manage unwanted populations.

The **German-Austrian** Black List Information System (GABLIS) has been developed as a generic risk assessment tool for invasive alien species in Germany and Austria, and is applicable to all groups of organisms. These assessments are not legally binding. The methodology has so far been tested for fish, vascular plants (including *Acer negundo*, *Ailanthus altissima*, *Fraxinus pennsylvanica*, *Paulownia tomentosa*, *Pinus nigra*, *P. strobus*, *Populus canadensis*, *Prunus laurocerasus*, *Prunus serotina*, *Pseudotsuga menziesii*, *Quercus rubra*, *Robinia pseudoacacia*), mammals, birds and macrozoobenthic species (Essl et al. 2011).

The Non-native Species Secretariat has responsibility for helping to coordinate the approach to invasive non-native species in **Great Britain**. Risk assessments are available for *Eucalyptus glaucescens* (low risk), *E. gunnii* (low risk), *E. nitens* (low risk) (<http://www.nonnativespecies.org/index.cfm?sectionid=51>).

In **Poland**, the use of *Ailanthus altissima* is banned, and could be permitted only by the General Directorate for Environment as stated in legislation (Dz.U. 2011 nr 210 poz. 1260)¹¹².

In **Slovakia** national legislation addresses the invasive alien species issue, e.g. in the Act No. 543/2002 Coll. on Nature and Landscape Protection as amended¹¹³. According to its provisions, it is prohibited to import, possess, grow, reproduce and trade invasive species and parts or products originating from them that could cause spontaneous dissemination of the invasive species. Moreover, land owners and land managers are obliged to eliminate invasive species from their land. According to the Order of the Ministry of Environment of SR No. 24/2003 Coll,¹¹⁴ Annex 2a, these provisions apply only to selected (the most problematic) invasive species. Two tree species, i.e. *Acer negundo* and *Ailanthus altissima* and two shrub species, i.e. *Amorpha fruticosa* and *Lycium barbarum* are listed. Information on invasive alien species (in the Slovak language) is available at: <http://www.sopsr.sk/publikacie/invazne/index.php>. In addition the national legislation in Slovakia provides also some general provisions for regulation of all "alien species" (non-native species not listed as the invasive ones). For instance according to § 7b of the above mentioned Act No 543/2002 Coll. only those wood alien species may be planted outside of built areas of villages - without permission of the competent state body - that are listed in the Annex 3a of the

¹¹¹ https://www.eppo.int/INVASIVE_PLANTS/ias_lists.htm

¹¹² *Rozporządzenie Ministra Środowiska z dnia 9 września 2011 r. w sprawie listy roślin i zwierząt gatunków obcych, które w przypadku uwolnienia do środowiska przyrodniczego mogą zagrozić gatunkom rodzimym lub siedliskom przyrodniczym* (<http://isap.sejm.gov.pl/DetailsServlet?id=WDU20112101260>). See also Tokarska-Guzik et al. (2012); Woziwoda et al. (2014).

¹¹³ <https://www.slov-lex.sk/pravne-predpisy/SK/ZZ/2002/543/20150101>

¹¹⁴ <https://www.slov-lex.sk/pravne-predpisy/SK/ZZ/2003/24/20150101>

above mentioned Order No. 24/2003 Coll. and in the Annex 1 of the Act No 138/2010 Coll. on Forest Reproduction Material¹¹⁵. Planting of other alien wood species outside of built areas of villages requires the permission of the competent state body. This permission is not required within the built areas. The Act No 138/2010 Coll. also specifies reforestation that is allowed again only by species listed in the Annex 1 of the Act.

In **Spain**, the Real Decreto 630/2013, “de 2 de agosto, por el que se regula el Catálogo español de especies exóticas invasoras” (Act 630/2013, 2nd August, that regulates Spanish Catalogue on Invasive Alien Species - <http://www.boe.es/boe/dias/2013/08/03/pdfs/BOE-A-2013-8565.pdf>) lists *Acacia dealbata*, *Acacia farnesiana*, *Acacia salicina* and *Ailanthus altissima*.

The **Swedish** forest legislation allows for regulations on the use of forest reproductive material in the establishment of new forest stands if warranted from a silvicultural point of view. Consequently, forest material from outside of the EU may not be introduced in Sweden without permit (Pettersson et al. 2016). In addition, “foreign” tree species may only be used as forest reproductive material in exceptional cases, although it is generally allowed to grow the *Pinus contorta* in certain parts of the country (Regulations from the Swedish Forest Agency, SKSFS 1993:2; SKSFS 2010:2.).

Black lists of invasive alien trees (which are not legally binding) have been published in many other countries, e.g. in **Romania** (Anastasiu & Negrean 2005).

Additional and updated information may be found either from national plant protection organisations (that is, Ministries of Agriculture¹¹⁶) or from Ministries of Environment in individual countries.

¹¹⁵ <https://www.slov-lex.sk/pravne-predpisy/SK/ZZ/2010/138/20140701>

¹¹⁶ Not in all the Countries. For example, the National Plant Protection Organization of the Netherlands was established in 1899. In 2012, the NPPO merged with other governmental organizations and formed the Netherlands Food and Consumer Product Safety Authority (NVWA). The NVWA is an integral part of the Ministry of Economic Affairs and its head office is based in Utrecht (<https://www.nvwa.nl/onderwerpen/english/dossier/national-plant-protection-organization-nppo>).